

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
27 January 2005 (27.01.2005)

PCT

(10) International Publication Number  
**WO 2005/007689 A1**

(51) International Patent Classification<sup>7</sup>: **C07K 14/045**,  
C12N 15/86, 7/04

Dr., Chapel Hill, NC 27517 (US). **MORRIS, John S.**  
[US/US]; 118 Trimble Ave., Cary, NC 27511 (US).

(21) International Application Number:  
PCT/US2004/021772

(74) Agent: **MYERS BIGEL SIBLEY & SAJOVEC, P.A.**;  
P.O. Box 37428, Raleigh, NC 27627 (US).

(22) International Filing Date: 8 July 2004 (08.07.2004)

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,  
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,  
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,  
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,  
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,  
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,  
ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/486,501 11 July 2003 (11.07.2003) US

(71) Applicant (for all designated States except US): **AL-  
PHAVAX, INC.** [US/US]; P.O. Box 110307, 2 Triangle  
Drive, Research Triangle Park, NC 27709-0307 (US).

(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,  
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,  
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,  
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG).


(72) Inventors; and

(75) Inventors/Applicants (for US only): **CHULAY, Jeffrey  
D.** [US/US]; 505 N. Boundary St., Chapel Hill, NC  
27514 (US). **DRYGA, Sergey** [US/US]; 211 Bonsail  
Place, Chapel Hill, NC 27514 (US). **REAP, Elizabeth  
A.** [US/US]; 12 Swallows Ridge Ct., Durham, NC 27713  
(US). **OLMSTED, Robert A.** [US/US]; 420 Bayberry

Published:  
— with international search report

[Continued on next page]

(54) Title: **ALPHAVIRUS-BASED CYTOMEGALOVIRUS VACCINES**



(57) Abstract: The present invention provides methods and compositions comprising a population of alphavirus replicon particles comprising alphavirus replicon RNAs, wherein a first replicon RNA comprises nucleic acid encoding cytomegalovirus pp65 and IE1 protein or immunogenic fragments thereof, and a second replicon RNA comprises nucleic acid encoding cytomegalovirus gB protein or an immunogenic fragment thereof, and wherein each of the two replicon RNAs is contained within a separate alphavirus replicon particle.

WO 2005/007689 A1

BEST AVAILABLE COPY



— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

## ALPHAVIRUS-BASED CYTOMEGALOVIRUS VACCINES

### RELATED APPLICATIONS

5           The present application claims the benefit, under 35 U.S.C. § 119(e), of U.S. provisional application serial number 60/486,501, filed July 11, 2003, the entire contents of which are incorporated by reference herein.

### BACKGROUND OF THE INVENTION

10

Human cytomegalovirus (HCMV) is a herpesvirus that causes widespread infection found across all geographic locations and socio-economic groups, with up to 85% of adults infected by age 40 in the United States. For most healthy people who acquire the virus after birth there are no long-term consequences. However, the risk of HCMV infection is  
15       significant for several high-risk groups including: (i) unborn children, (ii) adults who work with children, and (iii) immuno-compromised persons. The prevalence of these risk groups underlies the importance of the development of a safe and efficacious vaccine.

HCMV is typically secreted via a number of bodily fluids, e.g., saliva, urine and  
20       semen. Thus, transmission of the virus between people can occur through either sexual or non-sexual contact. An individual can contract HCMV through blood or organ transplants, and a mother can transmit it to her unborn fetus.

The virus demonstrates a life-long latency, but is most commonly non-symptomatic in  
25       healthy individuals. It sometimes can cause an illness with symptoms similar to those associated with mononucleosis. However, it can cause severe illness in immunocompromised individuals, e.g., transplant recipients or those with acquired immunodeficiency syndrome (AIDS), in addition to the severe, debilitating effects on unborn children whose immune systems have not yet matured.

30

In the case of transplants, bone marrow transplant recipients show a relatively high incidence of HCMV-induced pneumonia, with consequent high mortality among these patients. In solid organ transplant patients, disease triggered by HCMV can include a HCMV syndrome (consisting of fever and leucopenia), hepatitis, colitis and pneumonia. HCMV-  
35       induced disease in these transplant recipients is caused by the immunosuppressive effects of

the drugs required for transplant acceptance and the induction of graft vs. host disease (GVHD). The GVHD effect is most severe in those instances where the organ/marrow donor is HCMV seropositive and the recipient is HCMV seronegative.

- 5        For AIDS patients, HCMV is the most common opportunistic infection, in large part due to the fact that greater than 90% of HIV-infected individuals are co-infected with HCMV. In these patients, the infection most commonly manifests as retinitis, and usually occurs when the CD4+ cell counts are less than 50/ $\mu$ l. Prior to the adoption of highly active antiretroviral (HAART) protocols, 20-44% of AIDS patients developed HCMV disease.
- 10       While the use of HAART has also resulted in the reduction of HCMV disease, the unavailability of HAART for many AIDS patients, as well as the inability of many patients to tolerate HAART for extended periods of time, make the possibility of HCMV disease a continuing concern.

- 15       Congenital HCMV, a result of mother-to-fetus transmission, occurs at an overall rate of approximately 1%, but rates are much higher and symptomatic disease is more common when the mother has a primary infection. Women can be infected via sexual contact, since shedding of the virus from the cervix and in semen is common. Infected infants can remain viremic for up to five years after birth, becoming an important source for infection in day
- 20       care settings.

- Congenital HCMV can have horrific manifestations in infants. A fulminant cytomegalic inclusion disease can develop, characterized by jaundice, petechial rash, hepatosplenomegaly, microcephaly, and chorioretinitis. There is often progressive hearing
- 25       loss and mental retardation, which can be severe. The estimated costs to society in terms of care for victims of congenital HCMV are approximately four billion dollars.

- Thus, there remains a clear need for a safe and effective vaccine to combat HCMV infection, both prophylactically (for example, in adolescents or women of child-bearing
- 30       potential to prevent congenital infection or in HCMV-uninfected transplant candidates) and therapeutically (for example, in HCMV-infected transplant patients prior to and after transplantation of an organ or bone marrow).



## BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a diagram of the replicon vector VRP-pp65/IE1/Tr-gB.

5           Figure 2 shows the results of a CMV neutralization assay following immunization of mice with various CMV-VRP vaccines.

          Figures 3A-B show the results of an IFN- $\gamma$  ELISPOT assay following immunization of mice with CMV-VRP vaccines. Figure 3A shows the results obtained by using peptide  
10   pp65 #3 and a pp65 peptide pool to evaluate cellular immune responses. Figure 3B shows the results obtained by using peptide IE1 #50 and an IE1 peptide pool to evaluate cellular immune responses.

## SUMMARY OF THE INVENTION

15           The present invention provides a population of alphavirus replicon particles wherein said particles comprise alphavirus replicon RNAs, wherein a first replicon RNA comprises nucleic acid encoding cytomegalovirus pp65 and IE1 proteins or immunogenic fragments thereof, and a second replicon RNA comprises nucleic acid encoding cytomegalovirus gB  
20   protein or an immunogenic fragment thereof, and wherein each of the first and second replicon RNAs is contained within a separate alphavirus replicon particle.

          Further provided herein is a population of alphavirus replicon particles wherein said particles comprise a replicon RNA which comprises a regulatory cassette that directs  
25   transcription and translation of a nucleic acid encoding cytomegalovirus pp65 and IE1 proteins, or immunogenic fragments thereof.

          In additional embodiments, the present invention provides a population of alphavirus replicon particles wherein the particles comprise an alphavirus replicon RNA comprising  
30   nucleic acid encoding a CMV polypeptide selected from the group consisting of pp65, IE1, and gB, immunogenic fragments thereof or any combination thereof.

Also provided herein is a population of alphavirus replicon particles, wherein the particles comprise an alphavirus replicon RNA comprising nucleic acid encoding cytomegalovirus pp65 and gB proteins, or immunogenic fragments thereof.

5       The present invention also provides a population of alphavirus replicon particles comprising an alphavirus replicon RNA, wherein the replicon RNA of each particle comprises a first nucleic acid encoding cytomegalovirus pp65 protein or an immunogenic fragment thereof and a second nucleic acid encoding cytomegalovirus IE1 protein or an immunogenic fragment thereof, and wherein the expression of the first and second nucleic  
10       acid is controlled by separate regulatory cassettes.

Also provided herein are methods of inducing an immune response to CMV in a subject, comprising administering to the subject an effective amount of the populations of this invention.

15       Further provided is a method for inducing an immune response to CMV in a subject, comprising: a) priming the subject's immune system by administering to the subject an effective amount of a first immunizing component selected from the group consisting of: alphavirus replicon particles encoding CMV immunogens, CMV immunogens, nucleic acid  
20       molecules encoding CMV immunogens, a non-alphavirus viral vector encoding CMV immunogens, and any combination thereof; and b) boosting the subject's priming response by administering to the subject an effective amount of a second immunizing component selected from the group consisting of: alphavirus replicon particles encoding CMV immunogens, CMV immunogens, nucleic acid molecules encoding CMV immunogens, a  
25       non-alphavirus viral vector encoding CMV immunogens, and any combination thereof, wherein the first immunizing component is different from the second immunizing component and wherein at least the first immunizing component or the second immunizing component is an alphavirus replicon particle encoding CMV immunogens.

30

## DETAILED DESCRIPTION OF THE INVENTION

As used herein, "a," "an," or "the" can mean one or more than one. For example, "a cell" can mean one cell or a plurality of cells.

“Alphavirus” means a genus of viruses, all of which are members of the Togaviridae family. Known alphaviruses include Eastern Equine Encephalitis virus (EEE), Venezuelan Equine Encephalitis virus (VEE), Everglades virus, Mucambo virus, Pixuna virus, Western Equine Encephalitis virus (WEE), Sindbis virus, South African arbovirus 86 (S.A.AR86)  
5 Semliki Forest virus, Middleburg virus, Chikungunya virus, O’nyong-nyong virus, Ross River virus, Barmah Forest virus, Getah virus, Sagiya virus, Bebaru virus, Mayaro virus, Una virus, Aura virus, Whataroa virus, Babanki virus, Kyzylagach virus, Highlands J virus, Fort Morgan virus, Ndumu virus, and Buggy Creek virus. The alphaviral genome is a single-stranded, messenger-sense RNA, modified at the 5'-end with a methylated cap and at the 3'-  
10 end with a variable-length poly (A) tract. Structural subunits containing a single viral protein, capsid, associate with the RNA genome in an icosahedral nucleocapsid. In the virion, the capsid is surrounded by a lipid envelope covered with a regular array of transmembrane protein spikes, each of which consists of a heterodimeric complex of two glycoproteins, E1 and E2. See Pedersen et al., *J. Virol* 14:40 (1974). The Sindbis and  
15 Semliki Forest viruses are considered the prototypical alphaviruses and have been studied extensively. See Schlesinger, The Togaviridae and Flaviviridae, Plenum Publishing Corp., New York (1986). The preferred alphaviruses used in the constructs and methods of the claimed invention are VEE, S.AAR86, Sindbis (e.g., TR339, see U.S. Patent No. 6,008,035), and Semliki Forest Virus.

20

Within each named alphavirus, strains and/or subtypes are known. For example, several strains of the Venezuelan Equine Encephalitis virus (VEE) are known. Within the known strains of VEE, subtypes have been recognized. For example, the Trinidad Donkey strain is in subtype IA/B, and related subtypes include IC and IE. Virulent VEE strains have  
25 been isolated during mosquito-borne epizootic encephalomyelitis in equids in tropical and sub-tropical areas of the New World. The Trinidad Donkey strain is one of the virulent, epizootic strains, and it was passaged serially in tissue culture to create a live, attenuated strain (Berge et al. *Amer. J. Hyg.* 73:209-218 (1961)) known as TC-83. This strain, containing multiple attenuating mutations (see below, and Kinney et al. 1989 *Virology*  
30 170:19-30 (1989); with correction noted in Kinney et al. *J Virol* 67(3):1269-1277 (1993)) elicits VEE-specific neutralizing antibodies in most humans and equines and has been used successfully as a vaccine in both species (e.g., Pittman et al. *Vaccine* 14(4):337-343 (1996)). Thus, the TC-83 strain of VEE can also serve as the genetic background for an alphavirus replicon vector system as described herein.

The terms "alphavirus RNA replicon," "alphavirus replicon RNA," "alphavirus replicon vector" and "alphavirus RNA vector replicon" are used interchangeably to refer to an RNA molecule expressing nonstructural protein genes such that it can direct its own replication (amplification) and comprises, at a minimum, the 5' and 3' alphavirus replication recognition sequences, coding sequences for alphavirus nonstructural proteins, and a polyadenosine tract. It may additionally contain a regulatory cassette and a heterologous nucleic acid of interest that is expressed from the regulatory cassette. It may also be engineered to express one but not all alphavirus structural proteins.

10

Specific embodiments of the alphavirus RNA replicons utilized in the claimed invention may contain one or more "attenuating mutations," an attenuating mutation being a nucleotide deletion, addition, or substitution of one or more nucleotide(s), or a mutation that comprises rearrangement or chimeric construction which results in a loss of virulence in a live virus containing the mutation as compared to the appropriate wild-type alphavirus. Several examples of attenuating mutations have been previously described in U.S. Patent Nos. 5,639,650, 5,792,462 and 6,156,558. Specific attenuating mutations for the VEE E1 glycoprotein can include an attenuating mutation at any one of E1 amino acid positions 81, 272 and/or 253. Alphavirus particles made from the VEE-3042 mutant contain an isoleucine substitution at E1-81, and virus particles made from the VEE-3040 mutant contain an attenuating mutation at E1-253. Specific attenuating mutations for the VEE E2 glycoprotein can include an attenuating mutation at any one of E2 amino acid positions 76, 120, and/or 209. Alphavirus particles made from the VEE-3014 mutant contain attenuating mutations at both E1-272 and at E2-209 (see U.S. Patent No. 5,792,492). A specific attenuating mutation for the VEE E3 glycoprotein includes an attenuating mutation consisting of a deletion of E3 amino acids 56-59. Virus particles made from the VEE-3526 mutant, now being developed as a vaccine strain, contain this deletion in E3 (aa56-59) as well as a second attenuating mutation at E1-253.

30

Specific attenuating mutations for the S.A.AR86 E2 glycoprotein include an attenuating mutation at any one of E2 amino acid positions 304, 314, 372, and/or 376 (see U.S. Patent No. 5,639,650). Alternatively, the attenuating mutation can be a substitution, deletion and/or insertion of an amino acid in the E2 glycoprotein, for example, at any one or more of the following amino acid positions in any combination: 158, 159, 160, 161 and/or 162

(see Polo et al., PCT Publication No. WO 00/61772, the entire contents of which are incorporated by reference herein).

Attenuating mutations can also be present in the alphavirus non-structural proteins, nsp1-  
5 nsp4. Exemplary attenuating mutations in the non-structural proteins for S.A.AR86 include, but  
are not limited to, codons which specify an attenuating amino acid at any one or more of the  
following: nsp1 amino acid position 538, nsp2 amino acid position 96, nsp2 amino acid position  
372, nsp2 amino acid position 529; nsp2 amino acid position 571; nsp2 amino acid position 682;  
nsp2 amino acid position 804, nsp3 amino acid position 22, and in combination, codons at nsp2  
10 amino acid positions 529, 571, 682 and 804 and at nsp3 amino acid position. Other illustrative  
attenuating mutations for S.A.AR86 include those described in PCT Application No.  
PCT/US01/27644.

Another type of attenuating mutation of this invention can be one or more attenuating  
15 mutations in the non-translated regions of the alphavirus genome which cause a loss in virulence  
in a live virus containing such mutations (e.g., see Niesters and Strauss "Defined mutations in  
the 5' non-translated sequence of Sindbis virus RNA" *J Virol* 64: 4162-4168 (1990)). One  
example of such a mutation is at nucleotide 3 of the VEE genomic RNA, i.e., the third  
nucleotide following the 5' methylated cap (see, e.g., U.S. Patent No. 5,643,576, describing a  
20 G→C mutation at nt 3; and White et al. "Role of alpha/beta interferon in Venezuelan Equine  
Encephalitis virus pathogenesis: effect of an attenuating mutation in the 5' untranslated  
region" *J Virol* 75:2706-2718 (2000)). The mutation can be a G→A, U or C, but the G→A  
mutation is preferred for some embodiments.

25 The term "alphavirus structural protein/protein(s)" refers to one or a combination of  
the structural proteins encoded by an alphavirus. These are produced by the virus as a  
polyprotein and are represented generally in the literature as C-E3-E2-6k-E1. E3 and 6k  
serve as membrane translocation/transport signals for the two glycoproteins, E2 and E1.  
Thus, use of the term E1 herein can refer to E1, E3-E1, 6k-E1, or E3-6k-E1, and use of the  
30 term E2 herein can refer to E2, PE2, E3-E2, 6k-E2, or E3-6k-E2. As discussed above for the  
replicon, specific embodiments of the alphavirus structural proteins utilized in the claimed  
invention may contain one or more attenuating mutations, an attenuating mutation being a  
nucleotide deletion, addition, and/or substitution of one or more nucleotide(s), or a mutation

that comprises rearrangement or chimeric construction which results in a loss of virulence in a live virus containing the mutation as compared to the appropriate wild-type alphavirus.

The terms "alphavirus replicon particles (ARPs)," "virus replicon particles," and "recombinant alphavirus particles," used interchangeably herein, mean a virion-like structural complex incorporating an alphavirus replicon RNA that expresses one or more heterologous RNA sequences. Typically, the virion-like structural complex includes one or more alphavirus structural proteins embedded in a lipid envelope enclosing a nucleocapsid that in turn encloses the RNA. The lipid envelope is typically derived from the plasma membrane of the cell in which the particles are produced. Preferably, the alphavirus replicon RNA is surrounded by a nucleocapsid structure comprised of the alphavirus capsid protein, and the alphavirus glycoproteins are embedded in the cell-derived lipid envelope. ARPs are infectious but propagation-defective, i.e., the replicon RNA cannot propagate beyond the host cell into which the particles initially infect, in the absence of the helper nucleic acid(s) encoding the alphavirus structural proteins. The structural proteins and replicon RNA of the ARPs may be derived from the same or different alphaviruses. In one embodiment, the replicon RNA and the structural proteins are both derived from VEE, and such particles are sometimes referred to herein as "VRP" or "VRPs". In another embodiment, the replicon RNA is derived from VEE and the structural proteins are derived from Sindbis virus (see, e.g., Dubensky et al., U.S. Patent No. 6,376,236).

The term "helper(s)" refers to one or more nucleic acid molecules capable of being expressed to produce one or more alphavirus structural proteins. The helpers can be RNA or DNA molecules. In one embodiment, the helper is a single DNA molecule comprising a promoter capable of directing the expression of nucleic acid encoding all the structural proteins of the alphavirus. In another embodiment, the helper comprises two RNA molecules that together express nucleic acid encoding all the alphavirus structural proteins. These two RNA molecules can be produced *in vitro*, or they can be generated from a single DNA helper that resolves itself into two separate molecules *in vivo*. In the case of the DNA helper constructs that do not employ alphaviral recognition signals for replication and transcription, the theoretical frequency of recombination is lower than the bipartite RNA helper systems that employ such signals.

The terms "helper cell" and "packaging cell" are used interchangeably herein and refer to the cell in which alphavirus replicon particles are produced. The helper cell comprises a set of helpers that encode one or more alphavirus structural proteins. As disclosed herein, the helpers may be RNA or DNA. The cell can be any cell that is  
5 alphavirus-permissive, i.e., cells that are capable of producing alphavirus particles upon introduction of a viral RNA transcript. Alphavirus-permissive cells include, but are not limited to, Vero, baby hamster kidney (BHK), 293, 293T, chicken embryo fibroblast (CEF), and Chinese hamster ovary (CHO) cells. In certain embodiments of the claimed invention, the helper or packaging cell may additionally include a heterologous RNA-dependent RNA  
10 polymerase and/or a sequence-specific protease.

The term "immunogenic fragment" means a fragment (e.g., a peptide) of a CMV protein that can stimulate either humoral or cellular immune responses in the host.

15 To stimulate the humoral arm of the immune system, i.e., the production of antigen-specific antibodies, an immunogenic fragment can include at least about 5-10 contiguous amino acid residues of the full-length molecule, preferably at least about 15-25 contiguous amino acid residues of the full-length molecule, and most preferably at least about 20-50 or more contiguous amino acid residues of the full-length molecule, that define an epitope, or  
20 any integer between five amino acids and the full-length sequence, provided that the fragment in question retains immunogenic activity, as measured by any art-known assay, such as the ones described herein.

Regions of a given polypeptide that include an epitope can be identified using any  
25 number of epitope mapping techniques, well known in the art. (See, e.g., Epitope Mapping Protocols in Methods in Molecular Biology, Vol. 66, Glenn E. Morris, Ed., 1996, Humana Press, Totowa, N.J.) For example, linear epitopes can be determined by e.g., concurrently synthesizing large numbers of peptides on solid supports, the peptides corresponding to portions of the protein molecule, and reacting the peptides with antibodies while the peptides  
30 are still attached to the supports. Such techniques are known in the art and described in, e.g., U.S. Pat. No. 4,708,871; Geysen et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:3998-4002; Geysen et al. (1986) *Molec. Immunol.* 23:709-715, all incorporated herein by reference in their entireties.

Similarly, conformational epitopes are readily identified by determining spatial conformation of amino acids such as by, e.g., x-ray crystallography and 2-dimensional nuclear magnetic resonance. See, e.g., Epitope Mapping Protocols, supra. Antigenic regions of proteins can also be identified using standard antigenicity and hydropathy plots, such as  
5 those calculated using, e.g., the Omega version 1.0 software program available from the Oxford Molecular Group. This computer program employs the Hopp/Woods method (Hopp et al., *Proc. Natl. Acad. Sci USA* (1981) 78:3824-3828) for determining antigenicity profiles and the Kyte-Doolittle technique (Kyte et al., *J. Mol. Biol.* (1982) 157:105-132) for hydropathy plots.

10

Generally, T-cell epitopes that are involved in stimulating the cellular arm of a subject's immune system are short peptides of about 8-25 amino acids, and these are not typically predicted by the above-described methods for identifying humoral epitopes. A common way to identify T-cell epitopes is to use overlapping synthetic peptides and analyze  
15 pools of these peptides, or the individual ones, that are recognized by T cells from animals that are immune to the antigen of interest, using, for example, an enzyme-linked immunospot assay (ELISPOT). These overlapping peptides can also be used in other assays such as the stimulation of cytokine release or secretion, or evaluated by constructing major histocompatibility (MHC) tetramers containing the peptide. Such immunogenic fragments  
20 can also be identified based on their ability to stimulate lymphocyte proliferation in response to stimulation by various fragments from the antigen of interest.

25

The term "epitope" as used herein refers to a sequence of at least about 3 to 5, preferably about 5 to 10 or 15, and not more than about 1,000 amino acids (or any integer therebetween), which define a sequence that by itself or as part of a larger sequence, binds to an antibody generated in response to such sequence or stimulates a cellular immune response. There is no critical upper limit to the length of the fragment, which can comprise nearly the full-length of the protein sequence, or even a fusion protein comprising two or more epitopes from a single or multiple CMV proteins. An epitope for use in the subject invention is not  
30 limited to a polypeptide having the exact sequence of the portion of the parent protein from which it is derived. Indeed, there are many known strains or isolates of CMV and the virus retains the ability to continue to adapt, and there are several variable domains in the virus that exhibit relatively high degrees of variability between isolates. Thus the term "epitope" encompasses sequences identical to the native sequence, as well as modifications to the



native sequence, such as deletions, additions and substitutions (generally, but not always, conservative in nature).

The term "regulatory cassette" means a nucleic acid sequence encoding one or more elements necessary to direct transcription and/or translation of a nucleic acid encoding one or more polypeptides. In one embodiment, the regulatory cassette comprises only an alphavirus subgenomic promoter to direct transcription of a cytoplasmically located RNA that is then capped, and the capped end of the RNA directs translation of the subgenomic RNA. In another embodiment, the regulatory cassette comprises an internal ribosome entry site, or IRES, which directs translation of a downstream coding region. In another embodiment, the regulatory cassette comprises an alphavirus subgenomic promoter and an IRES, and it is engineered to allow the subgenomic promoter to direct transcription (and thus amplification) of the RNA sequence downstream from the promoter and to allow the IRES to direct translation of the subgenomic RNA.

15

"IRES" means an internal ribosome entry site. IRES sequences have been found in numerous transcripts from viruses that infect vertebrate and invertebrate cells as well as in transcripts from vertebrate and invertebrate genes. Examples of IRES elements suitable for use in this invention include: viral IRES elements from Picornaviruses e.g., poliovirus (PV), encephalomyocarditis virus (EMCV), foot-and-mouth disease virus (FMDV), from Flaviviruses e.g. hepatitis C virus (HCV), from Pestiviruses e.g., classical swine fever virus (CSFV), from Retroviruses e.g., murine leukemia virus (MLV), from Lentiviruses e.g., simian immunodeficiency virus (SIV), or cellular mRNA IRES elements such as those from translation initiation factors e.g., eIF4G or DAP5, from Transcription factors e.g., c-Myc (Yang and Sarnow, *Nucleic Acids Research* 25:2800-2807 1997) or NF- $\kappa$ B-repressing factor (NRF), from growth factors e.g., vascular endothelial growth factor (VEGF), fibroblast growth factor (FGF-2), platelet-derived growth factor B (PDGF B), from homeotic genes e.g., *Antennapedia*, from survival proteins e.g. X-Linked inhibitor of apoptosis (XIAP) or Apaf-1, or chaperones e.g. the immunoglobulin heavy-chain binding protein BiP (reviewed in Martínez-Salas et al., *Journal of General Virology* 82:973-984 (2001)).

25

30

Preferred IRES sequences that can be utilized in these embodiments are derived from: encephalomyocarditis virus (EMCV, ATCC accession # NC001479), cricket paralysis virus

(accession # AF218039), *Drosophila* C virus ATCC accession # AF014388, *Plautia stali* intestine virus (ATCC accession # AB006531), *Rhopalosiphum padi* virus (ATCC accession # AF022937), Himetobi P virus (ATCC accession # AB017037), acute bee paralysis virus (ATCC accession # AF150629), Black queen cell virus (ATCC accession # AF183905),  
5 Triatoma virus (ATCC accession # AF178440), *Acyrtosiphon pisum* virus (ATCC accession # AF024514), infectious flacherie virus (ATCC accession # AB000906), and Sacbrood virus (ATCC accession # AF092924). In addition to the naturally occurring IRES elements listed above, synthetic IRES sequences, designed to mimic the function of naturally occurring IRES sequences, can also be used. When more than one IRES is used in a replicon construct, the  
10 IRES elements may be the same or different.

"Boost" or "Booster" means a second immunization, after an initial (or "priming") immunization that enhances the immune response of the host. In one embodiment, the invention specifically provides a composition which produces an anamnestic response against  
15 a herpesvirus, e.g., CMV infection, in a sensitized subject, e.g., a horse, cow, or human, comprising an anamnestic response-inducing amount of a herpesvirus, e.g., CMV, immunizing component. As used herein, the term "anamnestic response" means a secondary (booster) immune response in a sensitized subject. By "sensitized subject" is meant a subject that has previously been in contact with herpesvirus, e.g., CMV, antigens either by natural  
20 exposure to the virus or by vaccination (primary immunization) with herpesvirus immunizing components, e.g., CMV-expressing alphavirus replicon particles.

At 230 kilobasepairs of double-stranded DNA, the cytomegalovirus genome is the largest  $\beta$ -herpesvirus known to infect humans. It has over 200 open reading frames  
25 responsible for encoding at least 165 genes; these are arranged in two segments, referred to as unique long ( $U_L$ ) and unique short ( $U_S$ ), which are separated by inverted repetitive nucleotide sequences. Thus, the choice of potential antigens to be used in a vaccine is quite large.

Some suggestions regarding vaccine approaches can be obtained from responses of  
30 healthy, seropositive individuals. In these individuals, 92% have CTLs present that target the pp65 antigen, 76% have CTLs to the IE1 antigen, 33% to the gB antigen, and 30% to the pp150 antigen (Gyulaj et al. 2000 *J. Infectious Diseases* 181:1537). In contrast, when this cell-mediated immunity is suppressed, the manifestations of HCMV disease are most severe. In addition, the reproductive number (number of cells infected by virus released from one

- infected cell) is reduced 2 to 7 times in HCMV-experienced hosts. Replication of HCMV in HCMV-experienced immuno-compromised hosts is delayed compared to HCMV-naïve hosts (doubling time 0.38 days and 1.12 days, respectively). As a corollary, cellular responses to pp65 and IE1 have also been demonstrated to protect from CMV infection in animal models
- 5 using the homologs of the HCMV genes encoding pp65 and IE1 (see Morello et al. *J. Virol.* 2000 Vol 74:3696). Finally, adoptive transfer of pp65-specific CTLs to bone-marrow transplant recipients protects them from CMV disease (Greenberg, P., Keystone Symposium April 2001; see also Walter et al., *N Engl J Med* 1995, 333:1038).
- 10       Antibodies to glycoprotein B (gB) are also present in HCMV infected individuals; these antibodies are neutralizing and have been implicated in protection of newborns from primary infection in animal models (Bourne et al. 2001 *J. Infectious Diseases* 183:59; Chatterjee et al. 2001 *J. Infectious Diseases* 183: 1547). Thus, it is expected that a preferred vaccine for humans provides both cellular and humoral immunity in order to be efficacious.
- 15 Live, attenuated vaccines using the Towne strain of HCMV have been used in transplant patients to reduce the severity of transplant-induced CMV in those seronegative recipients who received a seropositive kidney (Plotkin 1994 *Transplantation* 58:1176). However, because of concerns regarding the safety of live, attenuated vaccines derived from a virus that causes a chronic, persistent infection that can be transmitted to the fetus, and that can be
- 20 reactivated during periods of immunosuppression, alternative approaches are preferred.

- The alphavirus replicon vector system provides the opportunity to induce robust humoral and cellular immunity in humans. The replicon vector system is based on the replication machinery of an alphavirus, consisting of a replicon RNA vector and one or more
- 25 helper nucleic acids (reviewed in Rayner et al. (2002) *Rev. Med. Virol.* 12:279-96; see also U.S. Patent Nos. 5,792,462; 6,156,558; Pushko et al. (1997) *Virology* 239:389-401; U.S. Patent Publication No. 20020141975; PCT Publication No. WO 03/023026; the entire contents of which are incorporated herein by reference). The replicon RNA contains sequences required for replication and packaging of the RNA into a virus-like particle. It
- 30 expresses the nonstructural proteins required for genome replication and transcription of subgenomic RNA (if such constructs are utilized), but lacks the structural protein genes necessary for formation of viral particles. The replicon is engineered so that a regulatory cassette can direct the expression of a nucleic acid of interest, in this invention, one or more nucleic acids encoding CMV polypeptides or immunogenic fragments thereof. One or more

helper nucleic acids encode the alphavirus capsid and glycoproteins. When the replicon RNA vector and the one or more helper nucleic acids are introduced into an alphavirus-permissive cell, the replicon RNA is packaged into virus-like particles, which are harvested and purified to produce an immunogen, i.e., a vaccine composition.

5

In one embodiment, a replicon based on the Venezuelan Equine Encephalitis (VEE) virus is used as the vector for the CMV polypeptides. Nucleic acids encoding CMV proteins gB (e.g., from Towne strain), IE1 and pp65 (e.g., from AD169 strain) can be cloned into the alphavirus vector, e.g., the VEE vector, individually or in various combinations. Such combinations can encompass nucleic acid sequences encoding fusion proteins, e.g., IE1 and pp65 (or immunogenic fragments thereof) or they can encompass multiple regulatory cassettes within a single replicon, wherein each cassette drives either a single coding sequence or a nucleic acid sequence encoding a fusion protein of two or more CMV polypeptides, or immunogenic fragments thereof.

15

In one embodiment, each desired immunogen (i.e., a CMV polypeptide or an immunogenic fragment thereof) can be encoded by nucleic acid expressed in a separate alphavirus replicon particle, and mixtures of two or more ARPs, each comprising nucleic acid encoding a single CMV immunogen, can be prepared and administered to a subject. This invention therefore encompasses compositions comprising populations of one, two or three (or more) different ARPs. For example, in the case of three different ARPs, each ARP encodes a single CMV polypeptide, e.g., IE1, pp65, and gB (or immunogenic fragments thereof) and each ARP does not encode the other CMV polypeptides. In the case of two different ARPs, one ARP can express two CMV polypeptides, e.g., IE1 and pp65, either as fusion proteins or under the control of separate regulatory cassettes, as described above, and the second ARP can express one CMV polypeptide that is different, e.g., gB. Alternatively, a first ARP can express IE1 and a second ARP can express pp65. In another embodiment of a composition comprising two different ARPs, a first ARP can express pp65 and the second ARP can express gB. In an embodiment employing a single ARP, the replicon vector can include one regulatory cassette directing the expression of a nucleic acid encoding a fusion protein comprising at least pp65 and IE1, and in a specific embodiment, further comprising nucleic acid encoding gB. In another embodiment of the single ARP, one regulatory cassette can direct the expression of nucleic acid encoding gB, a second regulatory cassette can direct the expression of nucleic acid encoding IE1, and a third regulatory cassette can direct the

30

expression of nucleic acid encoding pp65. The regulatory cassette (RC)-encoding nucleic acid components can be in any order, e.g., RC-gB/RC-IE1/RC-pp65, and/or RC-pp65/RC-IE1/RC-gB. In another embodiment of the single ARP, a first regulatory cassette can direct the expression of a nucleic acid encoding a pp65-IE1 fusion protein, and a second RC can  
5 direct the expression of nucleic acid encoding gB. In all of the embodiments described herein, the use of the terms "pp65," "IE1," and "gB" are meant to encompass the full-length polypeptides, immunogenic fragments, and/or epitopes thereof. Nucleic acid sequences can be cloned from known CMV virus strains, e.g., Towne and AD169, and/or they can be synthetic sequences representing consensus sequences of the CMV sequences or epitopes  
10 from one or more CMV polypeptides, or chimeric sequences including fragments or epitopes from different CMV strains.

Thus, in certain embodiments, the present invention provides a population of alphavirus replicon particles wherein said particles comprise alphavirus replicon RNAs,  
15 wherein a first replicon RNA comprises nucleic acid encoding cytomegalovirus pp65 and IE1 proteins or immunogenic fragments thereof, and a second replicon RNA comprises nucleic acid encoding cytomegalovirus gB protein or an immunogenic fragment thereof, and wherein each of the first and second replicon RNAs is contained within a separate alphavirus replicon particle. In some embodiments, the first replicon RNA can direct transcription and translation  
20 of the nucleic acid encoding cytomegalovirus pp65 and IE1 proteins or immunogenic fragments thereof from two separate regulatory cassettes, whereby a first regulatory cassette directs transcription and translation of the CMV pp65-encoding nucleic acid and a second regulatory cassette directs replication and translation of the CMV IE1-encoding nucleic acid. In certain embodiments, the replicon RNAs of the particles of this population can further  
25 encode a CMV gB protein, and/or immunogenic fragment thereof and production of the CMV gB protein can be under the control of a separate, third regulatory cassette.

Further provided herein is a population of alphavirus replicon particles wherein the particles comprise a replicon RNA that comprises a regulatory cassette that directs  
30 transcription and translation of a nucleic acid encoding cytomegalovirus pp65 and IE1 proteins, or immunogenic fragments thereof.

Additionally provided herein is a population of alphavirus replicon particles wherein the particles comprise an alphavirus replicon RNA comprising nucleic acid encoding a CMV

polypeptide selected from the group consisting of pp65, IE1, and gB, immunogenic fragments thereof or any combination thereof.

5 In further embodiments, the present invention provides a population of alphavirus replicon particles, wherein the particles comprise an alphavirus replicon RNA comprising nucleic acid encoding-cytomegalovirus pp65 and gB proteins, or immunogenic fragments thereof. In some embodiments, this population can comprise alphavirus replicon RNA wherein the expression of the nucleic acid encoding cytomegalovirus pp65 and the expression of the nucleic acid encoding gB protein is controlled by separate regulatory cassettes.

10

A population of alphavirus replicon particles is also provided herein, comprising an alphavirus replicon RNA, wherein the replicon RNA of each particle comprises a first nucleic acid encoding cytomegalovirus pp65 protein or an immunogenic fragment thereof and a second nucleic acid encoding cytomegalovirus IE1 protein or an immunogenic fragment thereof, and wherein the expression of the first and second nucleic acid is controlled by separate regulatory cassettes.

15

In embodiments of this invention wherein the replicon RNAs direct expression of nucleic acid encoding CMV proteins and/or immunogenic fragments thereof of this invention from separate regulatory cassettes (e.g., one, two or three regulatory cassettes present on the same replicon RNA or on separate replicon RNAs of separate particles in a population), the regulatory cassettes can all be the same, the regulatory cassettes all be different and/or the regulatory cassettes can be present in any combination (e.g., two are the same and one is different).

20  
25

In some embodiments, the regulatory cassette of this invention can be an alphavirus subgenomic promoter. In other embodiments, the regulatory cassette of this invention can comprise (i) an alphavirus subgenomic promoter to direct transcription, and (ii) an IRES element to direct translation.

30

In those embodiments of this invention wherein nucleic acid encoding CMV pp65 protein and nucleic acid encoding CMV IE1 protein are present on the same replicon, the nucleic acid can be present as a coding sequence that produces a fusion protein of pp65 and IE1. A nonlimiting example of a nucleic acid encoding a pp65/IE1 fusion protein is provided

as SEQ ID NO:3 and a nonlimiting example of an amino acid sequence of a pp65/IE1 fusion protein is provided herein as SEQ ID NO:4. Other nucleic acids encoding the pp65/IE1 fusion protein of this invention would be readily determined by one of ordinary skill in the art and would vary based on the degeneracy of the DNA code. Other amino acid sequences  
5 having the functional characteristics of the pp65/IE1 fusion protein of this invention would be readily determined by one of ordinary skill in the art and would vary based on, for example conservative amino acid substitutions, as well as deletions and/or additions having a neutral or nominal effect on the functional characteristics of the fusion protein.

10 In those embodiments of this invention wherein the replicon RNA comprises nucleic acid encoding CMV gB protein or an immunogenic fragment thereof, the transmembrane domain of the gB protein or an immunogenic fragment thereof can be present or it can be deleted. A nonlimiting example of a nucleic acid encoding a CMV gB protein that has been truncated to delete the transmembrane domain is provided herein as SEQ ID NO:1. A  
15 nonlimiting example of an amino acid sequence of a truncated CMV gB protein is provided herein as SEQ ID NO:2. Other nucleic acids encoding the gB protein of this invention would be readily determined by one of ordinary skill in the art and would vary based on the degeneracy of the DNA code. Other amino acid sequences having the functional characteristics of the gB protein of this invention would be readily determined by one of  
20 ordinary skill in the art and would vary based on, for example conservative amino acid substitutions, as well as deletions and/or additions having a neutral or nominal effect on the functional characteristics of the gB protein.

Immunogenic fragments of the CMV proteins of this invention would be readily  
25 identified by one of ordinary skill in the art according to standard methods for identifying regions of immunogenicity in an amino acid sequence. Nonlimiting examples of immunogenic fragments of this invention are provided in the Sequence Listing included herewith and identified as SEQ ID NOs:5-262. These immunogenic fragments can be employed in any combination and in any ratio relative to one another in the compositions and  
30 methods of this invention. For example, "pools" of peptides can be created according to protocols standard in the art (see, e.g., Maecker et al. "Use of overlapping peptide mixtures as antigens for cytokine flow cytometry" *Journal of Immunological Methods* 255:27-40 (2001)) and used to evaluate the immune response in subjects infected with HCMV or immunized with HCMV vaccines to identify immunogenic fragments.

The CMV-expressing ARPs of this invention are formulated for use as pharmaceutical formulations, vaccines or immunogenic compositions, either for prophylaxis and/or treatment. These pharmaceutical formulations comprise a composition of this invention (e.g., infectious, propagation-defective ARPs) in combination with a pharmaceutically acceptable carrier.

Thus, in certain embodiments, the present invention provides a composition comprising an alphavirus particle of this invention in a pharmaceutically acceptable carrier. The compositions described herein can be formulated for administration in a pharmaceutical carrier in accordance with known techniques. See, e.g., Remington, *The Science And Practice of Pharmacy* (latest edition). In the manufacture of a pharmaceutical composition according to embodiments of the present invention, the composition of this invention is typically admixed with, *inter alia*, a pharmaceutically acceptable carrier. By "pharmaceutically acceptable carrier" is meant a carrier that is compatible with other ingredients in the pharmaceutical composition and that is not harmful or deleterious to the subject. The carrier may be a solid or a liquid, or both, and is preferably formulated with the composition of this invention as a unit-dose formulation. The pharmaceutical compositions are prepared by any of the well-known techniques of pharmacy including, but not limited to, admixing the components, optionally including one or more accessory ingredients. Exemplary pharmaceutically acceptable carriers include, but are not limited to, sterile pyrogen-free water and sterile pyrogen-free physiological saline solution. Such carriers can further include protein (e.g., serum albumin) and sugar (sucrose, sorbitol, glucose, etc.)

The pharmaceutical compositions of this invention include those suitable for oral, rectal, topical, inhalation (e.g., via an aerosol) buccal (e.g., sub-lingual), vaginal, parenteral (e.g., subcutaneous, intramuscular, intradermal, intraarticular, intrapleural, intraperitoneal, intracerebral, intraarterial, or intravenous), topical (i.e., both skin and mucosal surfaces, including airway surfaces) and transdermal administration. The compositions herein may also be administered via a skin scarification method, or transdermally via a patch or liquid. The compositions may be delivered subdermally in the form of a biodegradable material that releases the compositions over a period of time. The most suitable route in any given case will depend, as is well known in the art, on such factors as the species, age, gender and overall condition of the subject, the nature and severity of the condition being treated and/or



on the nature of the particular composition (i.e., dosage, formulation) that is being administered.

The ARPs can also be present in a formulation of this invention in an immunogenic amount. An "immunogenic amount" is an amount of the infectious alphavirus replicon particles which is sufficient to evoke an immune response in the subject to which the pharmaceutical formulation is administered. An amount of from about  $10^4$  to about  $10^{10}$ , preferably  $10^5$  to  $10^9$ , and in particular  $10^6$  to  $10^8$  infectious units (IU., as measured by indirect immunofluorescence assay), or ARPs, per dose can be administered to a subject, depending upon the age and species of the subject being treated.

Subjects to which effective and/or immunogenic amounts of the compositions of the present invention are administered include human and animal (e.g., mouse, monkey, guinea pig) subjects.

15

The vaccine compositions of this invention further comprise combinations of CMV polypeptide expressing ARPs with other CMV polypeptide expressing systems to provide the broadest (i.e., all aspects of the immune response, including those features described hereinabove) cellular and humoral responses possible. In certain embodiments, this can include the use of heterologous prime-boost strategies, in which the ARP compositions are used in combination with one or more of the following: recombinantly produced, purified CMV polypeptides (or immunogenic fragments thereof), naked nucleic acids encoding one or more CMV polypeptides, immunogenic fragments or epitopes, such nucleic acids formulated with lipid-containing moieties, non-alphavirus vectors (e.g., pox vectors, adenoviral vectors, herpes vectors, vesicular stomatitis virus vectors, paramyxoviral vectors, parvovirus vectors, papovavirus vectors, adeno-associated virus vectors and retroviral vectors) expressing one or more CMV immunogens, and other alphavirus vectors expressing one or more CMV immunogens. The viral vectors can be virus-like particles or nucleic acids. The alphavirus vectors can be replicon-containing particles, DNA-based replicon-containing vectors (sometimes referred to as an "ELVIS" system, see, for example, U.S. Patent No. 5,814,482) and/or naked RNA vectors.

30

Thus, the present invention further provides a method of inducing an immune response to CMV in a subject, comprising administering to the subject an effective amount of

the populations, particles and/or compositions of this invention. Also provided herein is a method of preventing or treating a CMV infection in a subject, comprising administering to the subject an effective amount of a population, particle and/or composition of this invention.

5           As used herein, an "effective amount" refers to an amount of a compound or composition that is sufficient to produce a desired effect, which can be a therapeutic, prophylactic and/or beneficial effect.

10           Also as used herein, the terms "treat," "treating" and "treatment" include any type of mechanism, action or activity that results in a change in the medical status of a subject, including an improvement in the condition of the subject (*e.g.*, change or improvement in one or more symptoms and/or clinical parameters), delay in the progression of the condition, prevention or delay of the onset of a disease or illness, etc.

15           In some embodiments, the present invention provides a method for inducing an immune response to CMV in a subject, comprising: a) priming the subject's immune system by administering to the subject an effective amount of a first immunizing component, which can be, but is not limited to, alphavirus replicon particles encoding CMV immunogens, CMV immunogens, nucleic acid molecules encoding CMV immunogens, a non-alphavirus viral  
20           vector encoding CMV immunogens, and any combination thereof; and b) boosting the subject's priming response by administering to the subject an effective amount of a second immunizing component, which can be, but is not limited to, alphavirus replicon particles encoding CMV immunogens, CMV immunogens, nucleic acid molecules encoding CMV immunogens, a non-alphavirus viral vector encoding CMV immunogens, and any  
25           combination thereof, wherein the first immunizing component can be different from the second immunizing component and wherein at least the first immunizing component or the second immunizing component is an alphavirus replicon particle encoding CMV immunogens.

30           In the methods of this invention, the first immunizing component can be a first alphavirus replicon particle and the second immunizing component can be a second alphavirus replicon particle, with the proviso that the first and second alphavirus particles are derived from different alphaviruses.

In other embodiments of the methods of this invention, the first immunizing component can comprise alphavirus replicon particles encoding cytomegalovirus pp65, IE1 and gB proteins or immunogenic fragments thereof, and the second immunizing component  
5 can comprise one or more CMV proteins and/or immunogenic fragments thereof.

In the methods of this invention, the immunizing components can be administered once or more than once (i.e., multiple times). For example, a first immunizing component of this invention and/or a second immunizing component of this invention can be administered  
10 one, two, three, four, five, six, seven, eight, nine or ten times at any time interval (e.g., hours, days, weeks, months, years, etc.) and in any of the amounts described herein, which can be the same amount each time or different amounts at different times of administration in any combination. In other embodiments, the administration of the first and second immunizing components can be combined or arranged in any order (e.g., the first and second immunizing  
15 components can be administered in an alternating sequence or in any other order).

In some embodiments of the present invention, the first and/or second immunizing component can be administered with an adjuvant. As used herein, "adjuvant" describes a substance, which can be any immunomodulating substance capable of being combined with  
20 the polypeptide or nucleic acid vaccine to enhance, improve or otherwise modulate an immune response in a subject without deleterious effect on the subject.

An adjuvant of this invention can be, but is not limited to, for example, an immunostimulatory cytokine (including, but not limited to, GM-CSF, interleukin-2, interleukin-12, interferon-gamma, interleukin-4, tumor necrosis factor-alpha, interleukin-1, hematopoietic factor flt3L, CD40L, B7.1 co-stimulatory molecules and B7.2 co-stimulatory  
25 molecules), SYNTEX adjuvant formulation 1 (SAF-1) composed of 5 percent (wt/vol) squalene (DASF, Parsippany, N.J.), 2.5 percent Pluronic, L121 polymer (Aldrich Chemical, Milwaukee), and 0.2 percent polysorbate (Tween 80, Sigma) in phosphate-buffered saline. Suitable adjuvants also include oil-in-water, saponin, an aluminum salt such as aluminum  
30 hydroxide gel (alum), aluminum phosphate, or alganmulin, but may also be a salt of calcium, iron or zinc, or may be an insoluble suspension of acylated tyrosine, or acylated sugars, cationically or anionically derivatized polysaccharides, or polyphosphazenes.

Other adjuvants are well known in the art and include QS-21, Freund's adjuvant (complete and incomplete), aluminum hydroxide, N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-normuramyl-L-alanyl-D-isoglutamine (CGP 11637, referred to as nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine (CGP 19835A, referred to as MTP-PE) and RIBI, which contains three components extracted from bacteria, monophosphoryl lipid A, trealose dimycolate and cell wall skeleton (MPL+TDM+CWS) in 2% squalene/Tween 80 emulsion.

Additional adjuvants can include, for example, a combination of monophosphoryl lipid A, preferably 3-de-O-acylated monophosphoryl lipid A (3D-MPL) together with an aluminum salt. An enhanced adjuvant system involves the combination of a monophosphoryl lipid A and a saponin derivative, particularly the combination of QS21 and 3D-MPL as disclosed in PCT publication number WO 94/00153 (the entire contents of which are incorporated herein by reference), or a less reactogenic composition where the QS21 is quenched with cholesterol as disclosed in PCT publication number WO 96/33739 (the entire contents of which are incorporated herein by reference). A particularly potent adjuvant formulation involving QS21 3D-MPL & tocopherol in an oil in water emulsion is described in PCT publication number WO 95/17210 (the entire contents of which are incorporated herein by reference). In addition, the nucleic acid of this invention can include a nucleotide sequence that provides an immunostimulatory signal and/or an adjuvant function, such as CpG sequences. Such CpG sequences, or motifs, are well known in the art.

An adjuvant of this invention, such as, for example, an immunostimulatory cytokine, can be administered before, concurrent with, and/or within a few hours, several hours, and/or 1, 2, 3, 4, 5, 6, 7, 8, 9, and/or 10 days before or after the administration of a composition of this invention to a subject.

Furthermore, any combination of adjuvants, such as immunostimulatory cytokines, can be co-administered to the subject before, after or concurrent with the administration of a composition of this invention. For example, combinations of immunostimulatory cytokines, can consist of two or more immunostimulatory cytokines of this invention, such as GM/CSF, interleukin-2, interleukin-12, interferon-gamma, interleukin-4, tumor necrosis factor-alpha,

interleukin-1, hematopoietic factor flt3L, CD40L, B7.1 co-stimulatory molecules and B7.2 co-stimulatory molecules. The effectiveness of an adjuvant or combination of adjuvants can be determined by measuring the immune response directed produced in response to administration of a composition of this invention to a subject with and without the adjuvant or combination of adjuvants, using standard procedures, as described herein and as known in the art.

In various embodiments of this invention comprising a non-alphavirus viral vector, the non-alphavirus viral vector can be, but is not limited to, a retroviral vector, an adenoviral vector, a poxvirus vector, a Vesicular Stomatitis Virus (VSV) vector or a picornavirus vector, as well as any other non-alphavirus viral vector now known or later identified.

The alphavirus particles employed in the methods of this invention can be particles derived from any alphavirus, such as, for example, Venezuelan Equine Encephalitis virus, S.A.AR86 virus, Semliki Forest virus, Sindbis virus, Ross River virus and any combination thereof. The alphavirus particles of this invention can also comprise elements (e.g., structural proteins/ replicon RNA) from two or more different alphaviruses to produce chimeric alphavirus particles (e.g., a particle comprising a Sindbis virus replicon RNA and VEE structural proteins). The production and testing of such chimeric particles is known in the art.

## EXAMPLES

### Example 1. Cloning of CMV Genes into VEE Replicon (Figure 1)

Standard molecular biology techniques were used in the cloning of all constructs and their analysis. The VEE replicon vector (Rayner et al.) was modified to introduce additional restriction sites for run-off transcription and ease of cloning. CMV genes coding for pp65 (UL83, strain AD169 (American Type Culture Collection No. VR-538), Immediate Early gene 1 (IE1, UL123, strain AD169), full-length glycoprotein B (gB, UL55, strain Towne (American Type Culture Collection No. VR-977) or C-terminus truncated gB (amino acids 1 – 692, excludes predicted transmembrane domain) were cloned under the control of a subgenomic ("SG") 26S promoter to generate replicons with single or multiple CMV genes.

In addition, a pp65-IE1 fusion construct was made, which was cloned into this modified VEE vector in a similar fashion. Table 1 identifies the various constructs that were made and tested. In some embodiments, a Csp45I restriction site was introduced into the replicon to linearize the vector, as a NotI site was identified in the pp65 coding sequence and NotI was used to linearize this vector for other uses, based on a NotI restriction site in the replicon sequence. In other embodiments, the pp65 coding sequence is modified to remove the NotI restriction site.

#### Example 2. Production of VEE Replicon Particles expressing CMV Genes

Packaged VEE Replicon Particles (VRP) were obtained after electroporation of CHO or Vero cells with *in vitro*-transcribed replicon and helper RNAs. Cells were maintained in EMEM (Vero) or F12-K (CHO) supplemented with 10% FBS in an atmosphere of 5% CO<sub>2</sub> at 37°C. For electroporation, cells were trypsinized and washed with phosphate buffered saline (PBS). Electroporation was performed using GenePulser Electroporator (Bio-Rad; Hercules, CA) and 0.4 cm cuvettes. After electroporation, the cells were resuspended in growth medium, seeded into tissue culture flasks containing growth medium, and incubated overnight. Growth medium containing released VRP was collected, filtered, and tested to confirm the absence of replication competent virus. VRP were then purified by affinity chromatography on HiTrap® heparin HP columns (Amersham, Piscataway, NJ), which are highly cross-linked agarose (6%), activated with N-hydroxysuccinimide and containing porcine heparin as the ligand. The VRP were formulated with 1% Human Serum Albumin and 5% sucrose in phosphate buffered saline.

#### Example 3. Protein Expression from CMV-expressing VRPs

Expression of nucleic acids to produce CMV proteins was analyzed by SDS-PAGE followed by silver stain (Invitrogen Inc., Carlsbad, CA) or western blot analysis with gene-specific monoclonal (Rumbaugh-Goodwin Institute) or goat polyclonal antibodies. Goat polyclonal antibodies were generated by immunization with purified CMV proteins.

Cells were infected with specified VRP at a multiplicity of infection (moi) of 10 IU/cell and incubated for 18-22 hr. Cell lysates in 0.5% SDS, 0.5% NP-40, 50 mM Tris-HCl,

pH 7.5, 0.1M NaCl, 1mM EDTA were normalized by protein content and 1 µg per lane of total protein was resolved on 4-12% gradient SDS-PAGE (Invitrogen Inc., Carlsbad, CA). Proteins were visualized by silver staining as recommended by manufacturer (Invitrogen Inc., Carlsbad, CA). Silver staining revealed prominent bands of the molecular weight expected for the CMV gene products.

Cells lysates prepared as described were analyzed by western blot with monoclonal or goat polyclonal monospecific antibodies specific for gB protein in reducing or non-reducing conditions. Western blot analysis revealed prominent bands of the molecular weight expected for the CMV gene products.

#### Example 4. Immunogenicity of VEE Replicon Particles expressing CMV Genes

Groups of 6-week-old female BALB/c mice (Charles River Laboratories, Raleigh, NC) were injected subcutaneously in both rear footpads with a total of  $10^6$  IU of VRP at weeks 0, 3, and 8. Serum samples were collected by retro-orbital bleed at day -1 (pre-bleed) and weeks 4 and 9. Spleens were harvested at week 15.

Some groups of 12 female BALB/c mice were primed and boosted with gB-VRP or truncated gB-VRP on Days 1 and 22. For the third inoculation given on Day 51, animals in these groups were split in half. Six of the animals in each group (Groups 3A and 4A) received a third VRP inoculation (the same VRP they received for the prime and boost) and the remaining six animals (Groups 3B and 4B) received truncated gB protein adjuvanted with RIBI adjuvant (Corixa Corporation, Seattle, WA) and bacterial endotoxin prepared in saline as their third inoculation. The VRPs were given at an inoculation dose of  $1 \times 10^6$  IU in the subcutaneous rear footpads. For the gB protein inoculations, 50 µg of protein in adjuvant was administered by intraperitoneal injection.

##### A. Induction of humoral immunity

A CMV neutralization assay was used to evaluate the humoral immune response to CMV gB. CMV neutralization titer was determined by incubating serial dilutions of heat-inactivated sera with a known concentration of CMV (Towne strain) in the presence of 5%

guinea pig complement (CedarLane Laboratories, Hornby, Ontario, Canada). Reduction of CMV infection was determined using a viral neutralization assay, as is standardly known in the art. Neutralization titer (NT<sub>50</sub>) was defined as 50% reduction in OD<sub>570</sub> compared to CMV-only control.

5

Immunization of mice with VRP expressing nucleic acid encoding glycoprotein B (full-length or truncated) resulted in induction of virus-neutralizing antibodies (Figure 2). Neutralizing antibody titers were significantly increased after the second boost. Higher neutralizing antibody titers were seen in Groups 3B and 4B with the addition of protein and  
10 adjuvant compared to the groups (Groups 3A and 4A) that got a third dose of VRP instead of protein and adjuvant (See Figure 2, Day 63).

#### B. Induction of cellular immunity

15

An IFN- $\gamma$  ELISPOT assay was used to evaluate the cellular immune response to pp65 and IE1. Splenic lymphocytes were prepared using Lympholyte M density gradient centrifugation after lysis of red blood cells. 96-well ELISPOT IP plates (Millipore, Bedford, MA) were coated with 1  $\mu$ g anti-mouse IFN- $\gamma$  mAb/well (MabTech, Mariemont, OH) and blocked with 10% FCS in RPMI-1640 (including supplements). 10<sup>6</sup> lymphocytes/well were  
20 plated alone or after mixing with Con A (4  $\mu$ g/ml) or peptide (10  $\mu$ g/ml). In all cases, each peptide was tested against known positive and negative lymphocytes and each lymphocyte preparation was tested against known positive and negative peptides. For detection, 0.1  $\mu$ g of biotinylated anti-mouse IFN- $\gamma$  (MabTech) was added to each well, followed by incubation with Avidin-Peroxidase Complex (Vector Laboratories, Burlingame, CA), and color  
25 development with AEC substrate. Spots were quantified by Zellnet, Inc. (New York, NY) using a Zeiss ELISPOT reader.

Cellular immune response of mice to CMV-expressing VRPs was measured by ELISPOT assay as described above. All constructs induced a robust immune response to  
30 pp65 and IE1 proteins (Figure 3).



**Example 5. Vaccine "Challenge" in Solid Organ Transplant Recipients**

A vaccine of the invention can be tested in "challenge" studies in humans undergoing solid organ transplant surgeries. CMV-seronegative patients on the organ/marrow waiting  
5 list are immunized, and then they undergo transplant surgery two weeks to several years after immunization. The "challenge" comes from the transplant itself, since most transplanted organs in the United States (> 60%) come from seropositive donors, and the CMV virus is transmitted via the organ. In addition, these CMV seronegative recipients of CMV seropositive organs are given booster doses of vaccine after the transplant surgery to maintain  
10 a sufficient level of immunity to prevent CMV disease. Booster doses are initially given at intervals of every one month to every six months. Patients are monitored and evaluated for at least one year, at monthly or quarterly intervals, and/or after the treatment for signs or symptoms of CMV disease.

**15 Example 6. Vaccination of Bone Marrow Transplant Recipients**

A vaccine of this invention can be administered to bone marrow transplant recipients to reduce or eliminate the transmission of HCMV via the donor bone marrow. The risk of disease from such HCMV transmission is particularly high in those seronegative recipients  
20 who receive a seropositive bone marrow. In one vaccination protocol, the bone marrow donor is vaccinated with a vaccine of the invention on one or more than one occasion, e.g., at six and two weeks before making the donation; the BMT recipient is vaccinated at intervals (e.g., every one to six months) starting about four weeks to about six months after receiving the transplanted bone marrow. The appearance of CMV viremia is monitored in the  
25 transplant recipient.

**Example 7. Vaccine "Challenge" in Pregnant Women**

A vaccine of the invention can be further tested in women of child-bearing potential  
30 who test seronegative for HCMV. Immunization protocols will typically include a priming immunization followed by one or two "booster" immunizations. These women are monitored for pregnancy outcomes, including the rates of CMV infection, symptomatic disease, and delayed sequelae in newborns.

Although the present process has been described with reference to specific details of certain embodiments thereof, it is not intended that such details should be regarded as limitations upon the scope of the invention except as and to the extent that they are included in the accompanying claims.

5

Throughout this application, various patents, patent publications and non-patent publications are referenced. The disclosures of these patents and publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention pertains.

Table 1. Organization of CMV genes in CMV-VRP vaccine candidates.

#	Designation	SG promoter #1	SG promoter #2	SG promoter #3
1	VRP-pp65	pp65		
2	VRP-IE1	IE1		
3	VRP-gB	gB		
4	VRP-Tr-gB	Tr-gB		
5	VRP-pp65/IE1	pp65	IE1	
9	VRP-pp65/gB	pp65	gB	
6	VRP-IE1/pp65	IE1	pp65	
11	VRP-pp65/IE1/gB	pp65	IE1	gB
10	VRP-pp65/IE1/Tr-gB	pp65	IE1	Tr-gB
7	VRP-pp65-IE1 fusion	pp65-IE1 fusion		
8	VRP-pp65-IE1 fusion/Tr-gB	pp65-IE1 fusion	Tr-gB	

What is claimed is:

1. A population of alphavirus replicon particles wherein said particles comprise alphavirus replicon RNAs, wherein a first replicon RNA comprises nucleic acid encoding cytomegalovirus pp65 and IE1 proteins or immunogenic fragments thereof, and a second replicon RNA comprises nucleic acid encoding cytomegalovirus gB protein or an immunogenic fragment thereof, and wherein each of the first and second replicon RNAs is contained within a separate alphavirus replicon particle.
2. A population of alphavirus replicon particles wherein said particles comprise a replicon RNA which comprises a regulatory cassette that directs transcription and translation of a nucleic acid encoding cytomegalovirus pp65 and IE1 proteins, or immunogenic fragments thereof.
3. The population of claim 2, wherein the nucleic acid encodes a fusion protein of pp65 and IE1.
4. The population of claim 1, wherein the first replicon RNA directs transcription and translation of the nucleic acid encoding cytomegalovirus pp65 and IE1 proteins or immunogenic fragments thereof from two separate regulatory cassettes, whereby a first regulatory cassette directs transcription and translation of the CMV pp65-encoding nucleic acid and a second regulatory cassette directs replication and translation of the CMV IE1-encoding nucleic acid.
5. The population of claim 1, wherein the first replicon RNA encodes a fusion protein of pp65 and IE1.
6. The population of claim 4, wherein the first and second regulatory cassettes are the same.
7. The population of claim 2, wherein the regulatory cassette is an alphavirus subgenomic promoter.

8. The population of claim 1 wherein the nucleic acid encoding cytomegalovirus gB protein or an immunogenic fragment thereof encodes a CMV gB protein or immunogenic fragment thereof wherein the transmembrane domain has been deleted.
9. The population of claim 4, wherein the first and second regulatory cassettes are different.
10. The population of claim 9, wherein the first and/or second regulatory cassette comprises (i) an alphavirus subgenomic promoter to direct transcription, and (ii) an IRES element to direct translation.
11. The population of claim 10 wherein the replicon RNA of each particle further encodes a CMV gB protein, or immunogenic fragment thereof and wherein production of the CMV gB protein is under the control of a separate, third regulatory cassette.
12. A population of alphavirus replicon particles wherein the particles comprise an alphavirus replicon RNA comprising nucleic acid encoding a CMV polypeptide selected from the group consisting of pp65, IE1, and gB, immunogenic fragments thereof or any combination thereof.
13. A population of alphavirus replicon particles, wherein the particles comprise an alphavirus replicon RNA comprising nucleic acid encoding-cytomegalovirus pp65 and gB proteins, or immunogenic fragments thereof.
14. The population of claim 13, wherein the expression of the nucleic acid encoding cytomegalovirus pp65 and the expression of the nucleic acid encoding gB protein is controlled by separate regulatory cassettes.
15. A population of alphavirus replicon particles comprising an alphavirus replicon RNA, wherein the replicon RNA of each particle comprises a first nucleic acid encoding cytomegalovirus pp65 protein or an immunogenic fragment thereof and a second nucleic acid encoding cytomegalovirus IE1 protein or an immunogenic fragment thereof, and wherein the expression of the first and second nucleic acid is controlled by separate regulatory cassettes.

16. A composition comprising the population of claim 1 in a pharmaceutically acceptable carrier.
17. A method of inducing an immune response to CMV in a subject, comprising administering to the subject an effective amount of the population of claim 1.
18. The method of claim 17, wherein the population is administered multiple times.
19. A method for inducing an immune response to CMV in a subject, comprising
  - a) priming the subject's immune system by administering to the subject an effective amount of a first immunizing component selected from the group consisting of:
    - alphavirus replicon particles encoding CMV immunogens,
    - CMV immunogens,
    - nucleic acid molecules encoding CMV immunogens,
    - a non-alphavirus viral vector encoding CMV immunogens, and
    - any combination thereof; and
  - b) boosting the subject's priming response by administering to the subject an effective amount of a second immunizing component selected from the group consisting of:
    - alphavirus replicon particles encoding CMV immunogens,
    - CMV immunogens,
    - nucleic acid molecules encoding CMV immunogens,
    - a non-alphavirus viral vector encoding CMV immunogens, and
    - any combination thereof,wherein the first immunizing component is different from the second immunizing component and wherein at least the first immunizing component or the second immunizing component is an alphavirus replicon particle encoding CMV immunogens.
20. The method of claim 19 wherein the first immunizing component is a first alphavirus replicon particle and the second immunizing component is a second alphavirus replicon particle, with the proviso that the first and second alphavirus particles are derived from different alphaviruses.
21. The method of claim 19 further comprising administering the first and/or second immunizing component multiple times.

22. The method of claim 19 wherein the first immunizing component comprises alphavirus replicon particles encoding cytomegalovirus pp65, IE1 and gB proteins or immunogenic fragments thereof, and wherein the second immunizing component comprises one or more CMV proteins and/or immunogenic fragments thereof.
23. The method of claim 22 wherein the CMV protein is a truncated gB protein.
24. The method of claim 19 wherein the first and/or second immunizing component is administered with an adjuvant.
25. The method of claim 24 wherein the adjuvant is selected from the group consisting of aluminum salts, oil-in-water, saponin, cytokines, oligonucleotides encoding immunostimulatory signals and any combination thereof.
26. The method of claim 19 wherein the non-alphavirus viral vector is selected from the group consisting of a retroviral vector, an adenoviral vector, a poxvirus vector, a Vesicular Stomatitis Virus (VSV) vector and a picornavirus vector.
27. The method of claim 19, wherein the alphavirus replicon particles are selected from the group consisting of particles derived from Venezuelan Equine Encephalitis virus, S.A.AR86 virus, Semliki Forest virus, Sindbis virus, Ross River virus and any combination thereof.
28. The method of claim 19, wherein the alphavirus replicon particles comprise elements from two or more alphaviruses.

Fig. 1









Fig. 2



Figs. 3A-B



## SEQUENCE LISTING

<110> Alphavax, Inc.  
 Chulay, Jeffrey D.  
 Dryga, Sergey A.  
 Reap, Elizabeth A.  
 Morris, John S.  
 Olmsted, Robert A.

<120> ALPHAVIRUS-BASED CYTOMEGALOVIRUS VACCINES

<130> 9368-7WO

<150> US 60/486,501

<151> 2003-07-11

<160> 262

<170> PatentIn version 3.2

<210> 1

<211> 2079

<212> DNA

<213> Human cytomegalovirus

<220>

<221> CDS

<222> (1)..(2076)

<223> Truncated gB

<400> 1

atg gaa tcc agg atc tgg tgc ctg gta gtc tgc gtt aac ttg tgt atc	48
Met Glu Ser Arg Ile Trp Cys Leu Val Val Cys Val Asn Leu Cys Ile	
1 5 10 15	

gtc tgt ctg ggt gct gcg gtt tcc tca tct tct act cgt gga act tct	96
Val Cys Leu Gly Ala Ala Val Ser Ser Ser Ser Thr Arg Gly Thr Ser	
20 25 30	

gct act cac agt cac cat tcc tct cat acg acg tct gct gct cat tct	144
Ala Thr His Ser His His Ser Ser His Thr Thr Ser Ala Ala His Ser	
35 40 45	

cga tcc ggt tca gtc tct caa cgc gta act tct tcc caa acg gtc agc	192
Arg Ser Gly Ser Val Ser Gln Arg Val Thr Ser Ser Gln Thr Val Ser	
50 55 60	

cat ggt gtt aac gag acc atc tac aac act acc ctc aag tac gga gat	240
His Gly Val Asn Glu Thr Ile Tyr Asn Thr Thr Leu Lys Tyr Gly Asp	
65 70 75 80	

gtg gtg ggg gtc aac acc acc aag tac ccc tat cgc gtg tgt tct atg	288
Val Val Gly Val Asn Thr Thr Lys Tyr Pro Tyr Arg Val Cys Ser Met	
85 90 95	

gca cag ggt acg gat ctt att cgc ttt gaa cgt aat atc gtc tgc acc	336
---	-----

Ala Gln Gly Thr Asp Leu Ile Arg Phe Glu Arg Asn Ile Val Cys Thr	
100 105 110	
tcg atg aag ccc atc aat gaa gac ctg gac gag ggc atc atg gtg gtc	384
Ser Met Lys Pro Ile Asn Glu Asp Leu Asp Glu Gly Ile Met Val Val	
115 120 125	
tac aaa cgc aac atc gtc gcg cac acc ttt aag gta cga gtc tac cag	432
Tyr Lys Arg Asn Ile Val Ala His Thr Phe Lys Val Arg Val Tyr Gln	
130 135 140	
aag gtt ttg acg ttt cgt cgt agc tac gct tac gtc cac acc act tat	480
Lys Val Leu Thr Phe Arg Arg Ser Tyr Ala Tyr Val His Thr Thr Tyr	
145 150 155 160	
ctg ctg ggc agc aac acg gaa tac gtg gcg cct cct atg tgg gag att	528
Leu Leu Gly Ser Asn Thr Glu Tyr Val Ala Pro Pro Met Trp Glu Ile	
165 170 175	
cat cat atc aac agt cac agt cag tgc tac agt tcc tac agc cgc gtt	576
His His Ile Asn Ser His Ser Gln Cys Tyr Ser Ser Tyr Ser Arg Val	
180 185 190	
ata gca ggc acg gtt ttc gtg gct tat cat agg gac agc tat gaa aac	624
Ile Ala Gly Thr Val Phe Val Ala Tyr His Arg Asp Ser Tyr Glu Asn	
195 200 205	
aaa acc atg caa tta atg ccc gac gat tat tcc aac acc cac agt acc	672
Lys Thr Met Gln Leu Met Pro Asp Asp Tyr Ser Asn Thr His Ser Thr	
210 215 220	
cgt tac gtg acg gtc aag gat caa tgg cac agc cgc ggc agc acc tgg	720
Arg Tyr Val Thr Val Lys Asp Gln Trp His Ser Arg Gly Ser Thr Trp	
225 230 235 240	
ctc tat cgt gag acc tgt aat ctg aat tgt atg gtg acc atc act act	768
Leu Tyr Arg Glu Thr Cys Asn Leu Asn Cys Met Val Thr Ile Thr Thr	
245 250 255	
gcg cgc tcc aag tat ccc tat cat ttt ttc gca act tcc acg ggt gat	816
Ala Arg Ser Lys Tyr Pro Tyr His Phe Phe Ala Thr Ser Thr Gly Asp	
260 265 270	
gtg gtt gac att tct cct ttc tac aac gga act aat cgc aat gcc agc	864
Val Val Asp Ile Ser Pro Phe Tyr Asn Gly Thr Asn Arg Asn Ala Ser	
275 280 285	
tat ttt gga gaa aac gcc gac aag ttt ttc att ttt ccg aac tac act	912
Tyr Phe Gly Glu Asn Ala Asp Lys Phe Phe Ile Phe Pro Asn Tyr Thr	
290 295 300	
atc gtc tcc gac ttt gga aga ccg aat tct gcg tta gag acc cac agg	960
Ile Val Ser Asp Phe Gly Arg Pro Asn Ser Ala Leu Glu Thr His Arg	
305 310 315 320	
ttg gtg gct ttt ctt gaa cgt gcg gac tca gtg atc tcc tgg gat ata	1008
Leu Val Ala Phe Leu Glu Arg Ala Asp Ser Val Ile Ser Trp Asp Ile	

325	330	335	
cag gac gag aag aat gtt act tgt caa ctc act ttc tgg gaa gcc tcg Gln Asp Glu Lys Asn Val Thr Cys Gln Leu Thr Phe Trp Glu Ala Ser 340	345	350	1056
gaa cgc acc att cgt tcc gaa gcc gag gac tcg tat cac ttt tct tct Glu Arg Thr Ile Arg Ser Glu Ala Glu Asp Ser Tyr His Phe Ser Ser 355	360	365	1104
gcc aaa atg acc gcc act ttc tta tct aag aag caa gag gtg aac atg Ala Lys Met Thr Ala Thr Phe Leu Ser Lys Lys Gln Glu Val Asn Met 370	375	380	1152
tcc gac tct gcg ctg gac tgt gta cgt gat gag gcc ata aat aag tta Ser Asp Ser Ala Leu Asp Cys Val Arg Asp Glu Ala Ile Asn Lys Leu 385	390	395	1200
cag cag att ttc aat act tca tac aat caa aca tat gaa aaa tat gga Gln Gln Ile Phe Asn Thr Ser Tyr Asn Gln Thr Tyr Glu Lys Tyr Gly 405	410	415	1248
aac gtg tcc gtc ttt gaa acc act ggt ggt ttg gtg gtg ttc tgg caa Asn Val Ser Val Phe Glu Thr Thr Gly Gly Leu Val Val Phe Trp Gln 420	425	430	1296
ggt atc aag caa aaa tct ctg gtg gaa ctc gaa cgt ttg gcc aac cgc Gly Ile Lys Gln Lys Ser Leu Val Glu Leu Glu Arg Leu Ala Asn Arg 435	440	445	1344
tcc agt ctg aat ctt act cat aat aga acc aaa aga agt aca gat ggc Ser Ser Leu Asn Leu Thr His Asn Arg Thr Lys Arg Ser Thr Asp Gly 450	455	460	1392
aac aat gca act cat tta tcc aac atg gag tcg gtg cac aat ctg gtc Asn Asn Ala Thr His Leu Ser Asn Met Glu Ser Val His Asn Leu Val 465	470	475	1440
tac gcc cag ctg cag ttc acc tat gac acg ttg cgc ggt tac atc aac Tyr Ala Gln Leu Gln Phe Thr Tyr Asp Thr Leu Arg Gly Tyr Ile Asn 485	490	495	1488
cgg gcg ctg gcg caa atc gca gaa gcc tgg tgt gtg gat caa cgg cgc Arg Ala Leu Ala Gln Ile Ala Glu Ala Trp Cys Val Asp Gln Arg Arg 500	505	510	1536
acc cta gag gtc ttc aag gaa ctt agc aag atc aac ccg tca gct att Thr Leu Glu Val Phe Lys Glu Leu Ser Lys Ile Asn Pro Ser Ala Ile 515	520	525	1584
ctc tcg gcc atc tac aac aaa ccg att gcc gcg cgt ttc atg ggc gat Leu Ser Ala Ile Tyr Asn Lys Pro Ile Ala Ala Arg Phe Met Gly Asp 530	535	540	1632
gtc ctg ggt ctg gcc agc tgc gtg acc att aac caa acc agc gtc aag Val Leu Gly Leu Ala Ser Cys Val Thr Ile Asn Gln Thr Ser Val Lys 545	550	555	1680
		560	

gtg ctg cgt gat atg aat gtg aag gaa tcg cca gga cgc tgc tac tca 1728  
 Val Leu Arg Asp Met Asn Val Lys Glu Ser Pro Gly Arg Cys Tyr Ser  
 565 570 575

cga cca gtg gtc atc ttt aat ttc gcc aac agc tcg tac gtg cag tac 1776  
 Arg Pro Val Val Ile Phe Asn Phe Ala Asn Ser Ser Tyr Val Gln Tyr  
 580 585 590

ggc caa ctg ggc gag gat aac gaa atc ctg ttg ggc aac cac cgc act 1824  
 Gly Gln Leu Gly Glu Asp Asn Glu Ile Leu Leu Gly Asn His Arg Thr  
 595 600 605

gag gaa tgt cag ctt ccc agc ctc aag atc ttc atc gcc ggc aac tcg 1872  
 Glu Glu Cys Gln Leu Pro Ser Leu Lys Ile Phe Ile Ala Gly Asn Ser  
 610 615 620

gcc tac gag tac gtg gac tac ctc ttc aaa cgc atg att gac ctc agc 1920  
 Ala Tyr Glu Tyr Val Asp Tyr Leu Phe Lys Arg Met Ile Asp Leu Ser  
 625 630 635 640

agc atc tcc acc gtc gac agc atg atc gcc cta gac atc gac ccg ctg 1968  
 Ser Ile Ser Thr Val Asp Ser Met Ile Ala Leu Asp Ile Asp Pro Leu  
 645 650 655

gaa aac acc gac ttc agg gta ctg gaa ctt tac tcg cag aaa gaa ttg 2016  
 Glu Asn Thr Asp Phe Arg Val Leu Glu Leu Tyr Ser Gln Lys Glu Leu  
 660 665 670

cgt tcc agc aac gtt ttt gat ctc gag gag atc atg cgc gag ttc aat 2064  
 Arg Ser Ser Asn Val Phe Asp Leu Glu Glu Ile Met Arg Glu Phe Asn  
 675 680 685

tcg tat aag cag tga 2079  
 Ser Tyr Lys Gln  
 690

<210> 2

<211> 692

<212> PRT

<213> Human cytomegalovirus

<400> 2

Met Glu Ser Arg Ile Trp Cys Leu Val Val Cys Val Asn Leu Cys Ile  
 1 5 10 15

Val Cys Leu Gly Ala Ala Val Ser Ser Ser Ser Thr Arg Gly Thr Ser  
 20 25 30

Ala Thr His Ser His His Ser Ser His Thr Thr Ser Ala Ala His Ser  
 35 40 45

Arg Ser Gly Ser Val Ser Gln Arg Val Thr Ser Ser Gln Thr Val Ser  
50 55 60

His Gly Val Asn Glu Thr Ile Tyr Asn Thr Thr Leu Lys Tyr Gly Asp  
65 70 75 80

Val Val Gly Val Asn Thr Thr Lys Tyr Pro Tyr Arg Val Cys Ser Met  
85 90 95

Ala Gln Gly Thr Asp Leu Ile Arg Phe Glu Arg Asn Ile Val Cys Thr  
100 105 110

Ser Met Lys Pro Ile Asn Glu Asp Leu Asp Glu Gly Ile Met Val Val  
115 120 125

Tyr Lys Arg Asn Ile Val Ala His Thr Phe Lys Val Arg Val Tyr Gln  
130 135 140

Lys Val Leu Thr Phe Arg Arg Ser Tyr Ala Tyr Val His Thr Thr Tyr  
145 150 155 160

Leu Leu Gly Ser Asn Thr Glu Tyr Val Ala Pro Pro Met Trp Glu Ile  
165 170 175

His His Ile Asn Ser His Ser Gln Cys Tyr Ser Ser Tyr Ser Arg Val  
180 185 190

Ile Ala Gly Thr Val Phe Val Ala Tyr His Arg Asp Ser Tyr Glu Asn  
195 200 205

Lys Thr Met Gln Leu Met Pro Asp Asp Tyr Ser Asn Thr His Ser Thr  
210 215 220

Arg Tyr Val Thr Val Lys Asp Gln Trp His Ser Arg Gly Ser Thr Trp  
225 230 235 240

Leu Tyr Arg Glu Thr Cys Asn Leu Asn Cys Met Val Thr Ile Thr Thr  
245 250 255

Ala Arg Ser Lys Tyr Pro Tyr His Phe Phe Ala Thr Ser Thr Gly Asp  
260 265 270

Val Val Asp Ile Ser Pro Phe Tyr Asn Gly Thr Asn Arg Asn Ala Ser

275	280	285
Tyr Phe Gly Glu Asn Ala Asp Lys Phe Phe Ile Phe Pro Asn Tyr Thr 290 295 300		
Ile Val Ser Asp Phe Gly Arg Pro Asn Ser Ala Leu Glu Thr His Arg 305 310 315 320		
Leu Val Ala Phe Leu Glu Arg Ala Asp Ser Val Ile Ser Trp Asp Ile 325 330 335		
Gln Asp Glu Lys Asn Val Thr Cys Gln Leu Thr Phe Trp Glu Ala Ser 340 345 350		
Glu Arg Thr Ile Arg Ser Glu Ala Glu Asp Ser Tyr His Phe Ser Ser 355 360 365		
Ala Lys Met Thr Ala Thr Phe Leu Ser Lys Lys Gln Glu Val Asn Met 370 375 380		
Ser Asp Ser Ala Leu Asp Cys Val Arg Asp Glu Ala Ile Asn Lys Leu 385 390 395 400		
Gln Gln Ile Phe Asn Thr Ser Tyr Asn Gln Thr Tyr Glu Lys Tyr Gly 405 410 415		
Asn Val Ser Val Phe Glu Thr Thr Gly Gly Leu Val Val Phe Trp Gln 420 425 430		
Gly Ile Lys Gln Lys Ser Leu Val Glu Leu Glu Arg Leu Ala Asn Arg 435 440 445		
Ser Ser Leu Asn Leu Thr His Asn Arg Thr Lys Arg Ser Thr Asp Gly 450 455 460		
Asn Asn Ala Thr His Leu Ser Asn Met Glu Ser Val His Asn Leu Val 465 470 475 480		
Tyr Ala Gln Leu Gln Phe Thr Tyr Asp Thr Leu Arg Gly Tyr Ile Asn 485 490 495		
Arg Ala Leu Ala Gln Ile Ala Glu Ala Trp Cys Val Asp Gln Arg Arg 500 505 510		



Thr Leu Glu Val Phe Lys Glu Leu Ser Lys Ile Asn Pro Ser Ala Ile  
 515 520 525

Leu Ser Ala Ile Tyr Asn Lys Pro Ile Ala Ala Arg Phe Met Gly Asp  
 530 535 540

Val Leu Gly Leu Ala Ser Cys Val Thr Ile Asn Gln Thr Ser Val Lys  
 545 550 555 560

Val Leu Arg Asp Met Asn Val Lys Glu Ser Pro Gly Arg Cys Tyr Ser  
 565 570 575

Arg Pro Val Val Ile Phe Asn Phe Ala Asn Ser Ser Tyr Val Gln Tyr  
 580 585 590

Gly Gln Leu Gly Glu Asp Asn Glu Ile Leu Leu Gly Asn His Arg Thr  
 595 600 605

Glu Glu Cys Gln Leu Pro Ser Leu Lys Ile Phe Ile Ala Gly Asn Ser  
 610 615 620

Ala Tyr Glu Tyr Val Asp Tyr Leu Phe Lys Arg Met Ile Asp Leu Ser  
 625 630 635 640

Ser Ile Ser Thr Val Asp Ser Met Ile Ala Leu Asp Ile Asp Pro Leu  
 645 650 655

Glu Asn Thr Asp Phe Arg Val Leu Glu Leu Tyr Ser Gln Lys Glu Leu  
 660 665 670

Arg Ser Ser Asn Val Phe Asp Leu Glu Glu Ile Met Arg Glu Phe Asn  
 675 680 685

Ser Tyr Lys Gln  
 690

<210> 3  
 <211> 3156  
 <212> DNA  
 <213> Artificial

<220>  
 <223> hCMV pp65-IE1 fusion

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(3153)

&lt;400&gt; 3

atg gag tcg cgc ggt cgc cgt tgt ccc gaa atg ata tcc gta ctg ggt	48
Met Glu Ser Arg Gly Arg Arg Cys Pro Glu Met Ile Ser Val Leu Gly	
1 5 10 15	
ccc att tcg ggg cac gtg ctg aaa gcc gtg ttt agt cgc ggc gat acg	96
Pro Ile Ser Gly His Val Leu Lys Ala Val Phe Ser Arg Gly Asp Thr	
20 25 30	
ccg gtg ctg ccg cac gag acg cga ctc ctg cag acg ggt atc cac gta	144
Pro Val Leu Pro His Glu Thr Arg Leu Leu Gln Thr Gly Ile His Val	
35 40 45	
cgc gtg agc cag ccc tcg ctg atc ttg gta tcg cag tac acg ccc gac	192
Arg Val Ser Gln Pro Ser Leu Ile Leu Val Ser Gln Tyr Thr Pro Asp	
50 55 60	
tcg acg cca tgc cac cgc ggc gac aat cag ctg cag gtg cag cac acg	240
Ser Thr Pro Cys His Arg Gly Asp Asn Gln Leu Gln Val Gln His Thr	
65 70 75 80	
tac ttt acg ggc agc gag gtg gag aac gtg tcg gtc aac gtg cac aac	288
Tyr Phe Thr Gly Ser Glu Val Glu Asn Val Ser Val Asn Val His Asn	
85 90 95	
ccc acg ggc cga agc atc tgc ccc agc cag gag ccc atg tcg atc tat	336
Pro Thr Gly Arg Ser Ile Cys Pro Ser Gln Glu Pro Met Ser Ile Tyr	
100 105 110	
gtg tac gcg ctg ccg ctc aag atg ctg aac atc ccc agc atc aac gtg	384
Val Tyr Ala Leu Pro Leu Lys Met Leu Asn Ile Pro Ser Ile Asn Val	
115 120 125	
cac cac tac ccg tcg gcg gcc gag cgc aaa cac cga cac ctg ccc gta	432
His His Tyr Pro Ser Ala Ala Glu Arg Lys His Arg His Leu Pro Val	
130 135 140	
gct gac gct gtg att cac gcg tcg ggc aag cag atg tgg cag gcg cgt	480
Ala Asp Ala Val Ile His Ala Ser Gly Lys Gln Met Trp Gln Ala Arg	
145 150 155 160	
ctc acg gtc tcg gga ctg gcc tgg acg cgt cag cag aac cag tgg aaa	528
Leu Thr Val Ser Gly Leu Ala Trp Thr Arg Gln Gln Asn Gln Trp Lys	
165 170 175	
gag ccc gac gtc tac tac acg tca gcg ttc gtg ttt ccc acc aag gac	576
Glu Pro Asp Val Tyr Tyr Thr Ser Ala Phe Val Phe Pro Thr Lys Asp	
180 185 190	
gtg gca ctg cgg cac gtg gtg tgc gcg cac gag ctg gtt tgc tcc atg	624
Val Ala Leu Arg His Val Val Cys Ala His Glu Leu Val Cys Ser Met	

195	200	205	
gag aac acg cgc gca acc aag atg cag gtg ata ggt gac cag tac gtc Glu Asn Thr Arg Ala Thr Lys Met Gln Val Ile Gly Asp Gln Tyr Val 210 215 220			672
aag gtg tac ctg gag tcc ttc tgc gag gac gtg ccc tcc ggc aag ctc Lys Val Tyr Leu Glu Ser Phe Cys Glu Asp Val Pro Ser Gly Lys Leu 225 230 235 240			720
ttt atg cac gtc acg ctg ggc tct gac gtg gaa gag gac ctg acg atg Phe Met His Val Thr Leu Gly Ser Asp Val Glu Glu Asp Leu Thr Met 245 250 255			768
acc cgc aac ccg caa ccc ttc atg cgc ccc cac gag cgc aac ggc ttt Thr Arg Asn Pro Gln Pro Phe Met Arg Pro His Glu Arg Asn Gly Phe 260 265 270			816
acg gtg ttg tgt ccc aaa aat atg ata atc aaa ccg ggc aag atc tcg Thr Val Leu Cys Pro Lys Asn Met Ile Ile Lys Pro Gly Lys Ile Ser 275 280 285			864
cac atc atg ctg gat gtg gct ttt acc tca cac gag cat ttt ggg ctg His Ile Met Leu Asp Val Ala Phe Thr Ser His Glu His Phe Gly Leu 290 295 300			912
ctg tgt ccc aag agc atc ccg ggc ctg agc atc tca ggt aac ctg ttg Leu Cys Pro Lys Ser Ile Pro Gly Leu Ser Ile Ser Gly Asn Leu Leu 305 310 315 320			960
atg aac ggg cag cag atc ttc ctg gag gta caa gcc ata cgc gag acc Met Asn Gly Gln Gln Ile Phe Leu Glu Val Gln Ala Ile Arg Glu Thr 325 330 335			1008
gtg gaa ctg cgt cag tac gat ccc gtg gct gcg ctc ttc ttt ttc gat Val Glu Leu Arg Gln Tyr Asp Pro Val Ala Ala Leu Phe Phe Phe Asp 340 345 350			1056
atc gac ttg ctg ctg cag cgc ggg cct cag tac agc gag cac ccc acc Ile Asp Leu Leu Leu Gln Arg Gly Pro Gln Tyr Ser Glu His Pro Thr 355 360 365			1104
ttc acc agc cag tat cgc atc cag ggc aag ctt gag tac cga cac acc Phe Thr Ser Gln Tyr Arg Ile Gln Gly Lys Leu Glu Tyr Arg His Thr 370 375 380			1152
tgg gac cgg cac gac gag ggt gcc gcc cag ggc gac gac gac gtc tgg Trp Asp Arg His Asp Glu Gly Ala Ala Gln Gly Asp Asp Asp Val Trp 385 390 395 400			1200
acc agc gga tcg gac tcc gac gaa gaa ctc gta acc acc gag cgc aag Thr Ser Gly Ser Asp Ser Asp Glu Glu Leu Val Thr Thr Glu Arg Lys 405 410 415			1248
acg ccc cgc gtc acc ggc ggc ggc gcc atg gcg ggc gcc tcc act tcc Thr Pro Arg Val Thr Gly Gly Gly Ala Met Ala Gly Ala Ser Thr Ser 420 425 430			1296

gcg ggc cgc aaa cgc aaa tca gca tcc tcg gcg acg gcg tgc acg tcg Ala Gly Arg Lys Arg Lys Ser Ala Ser Ser Ala Thr Ala Cys Thr Ser 435 440 445	1344
ggc gtt atg aca cgc ggc cgc ctt aag gcc gag tcc acc gtc gcg ccc Gly Val Met Thr Arg Gly Arg Leu Lys Ala Glu Ser Thr Val Ala Pro 450 455 460	1392
gaa gag gac acc gac gag gat tcc gac aac gaa atc cac aat ccg gcc Glu Glu Asp Thr Asp Glu Asp Ser Asp Asn Glu Ile His Asn Pro Ala 465 470 475 480	1440
gtg ttc acc tgg ccg ccc tgg cag gcc ggc atc ctg gcc cgc aac ctg Val Phe Thr Trp Pro Pro Trp Gln Ala Gly Ile Leu Ala Arg Asn Leu 485 490 495	1488
gtg ccc atg gtg gct acg gtt cag ggt cag aat ctg aag tac cag gaa Val Pro Met Val Ala Thr Val Gln Gly Gln Asn Leu Lys Tyr Gln Glu 500 505 510	1536
ttc ttc tgg gac gcc aac gac atc tac cgc atc ttc gcc gaa ttg gaa Phe Phe Trp Asp Ala Asn Asp Ile Tyr Arg Ile Phe Ala Glu Leu Glu 515 520 525	1584
ggc gta tgg cag ccc gct gcg caa ccc aaa cgt cgc cgc cac cgg caa Gly Val Trp Gln Pro Ala Ala Gln Pro Lys Arg Arg Arg His Arg Gln 530 535 540	1632
gac gcc ttg ccc ggg cca tgc atc gcc tcg acg ccc aaa aag cac cga Asp Ala Leu Pro Gly Pro Cys Ile Ala Ser Thr Pro Lys Lys His Arg 545 550 555 560	1680
ggc gag tcc tct gcc aag aga aag atg gac cct gat aat cct gac gag Gly Glu Ser Ser Ala Lys Arg Lys Met Asp Pro Asp Asn Pro Asp Glu 565 570 575	1728
ggc cct tcc tcc aag gtg cca cgg ccc gag aca ccc gtg acc aag gcc Gly Pro Ser Ser Lys Val Pro Arg Pro Glu Thr Pro Val Thr Lys Ala 580 585 590	1776
acg acg ttc ctg cag act atg ttg agg aag gag gtt aac agt cag ctg Thr Thr Phe Leu Gln Thr Met Leu Arg Lys Glu Val Asn Ser Gln Leu 595 600 605	1824
agt ctg gga gac ccg ctg ttt cca gag ttg gcc gaa gaa tcc ctc aaa Ser Leu Gly Asp Pro Leu Phe Pro Glu Leu Ala Glu Glu Ser Leu Lys 610 615 620	1872
act ttt gaa caa gtg acc gag gat tgc aac gag aac ccc gag aaa gat Thr Phe Glu Gln Val Thr Glu Asp Cys Asn Glu Asn Pro Glu Lys Asp 625 630 635 640	1920
gtc ctg gca gaa ctc gtc aaa cag att aag gtt cga gtg gac atg gtg Val Leu Ala Glu Leu Val Lys Gln Ile Lys Val Arg Val Asp Met Val 645 650 655	1968

cgg cat aga atc aag gag cac atg ctg aaa aaa tat acc cag acg gaa Arg His Arg Ile Lys Glu His Met Leu Lys Lys Tyr Thr Gln Thr Glu 660 665 670	2016
gag aaa ttc act ggc gcc ttt aat atg atg gga gga tgt ttg cag aat Glu Lys Phe Thr Gly Ala Phe Asn Met Met Gly Gly Cys Leu Gln Asn 675 680 685	2064
gcc tta gat atc tta gat aag gtt cat gag cct ttc gag gag atg aag Ala Leu Asp Ile Leu Asp Lys Val His Glu Pro Phe Glu Glu Met Lys 690 695 700	2112
tgt att ggg cta act atg cag agc atg tat gag aac tac att gta cct Cys Ile Gly Leu Thr Met Gln Ser Met Tyr Glu Asn Tyr Ile Val Pro 705 710 715 720	2160
gag gat aag cgg gag atg tgg atg gct tgt att aag gag ctg cat gat Glu Asp Lys Arg Glu Met Trp Met Ala Cys Ile Lys Glu Leu His Asp 725 730 735	2208
gtg agc aag ggc gcc gct aac aag ttg ggg ggt gca ctg cag gct aag Val Ser Lys Gly Ala Ala Asn Lys Leu Gly Gly Ala Leu Gln Ala Lys 740 745 750	2256
gcc cgt gct aaa aag gat gaa ctt agg aga aag atg atg tat atg tgc Ala Arg Ala Lys Lys Asp Glu Leu Arg Arg Lys Met Met Tyr Met Cys 755 760 765	2304
tac agg aat ata gag ttc ttt acc aag aac tca gcc ttc cct aag acc Tyr Arg Asn Ile Glu Phe Phe Thr Lys Asn Ser Ala Phe Pro Lys Thr 770 775 780	2352
acc aat ggc tgc agt cag gcc atg gcg gca ctg cag aac ttg cct cag Thr Asn Gly Cys Ser Gln Ala Met Ala Ala Leu Gln Asn Leu Pro Gln 785 790 795 800	2400
tgc tcc cct gat gag att atg gct tat gcc cag aaa ata ttt aag att Cys Ser Pro Asp Glu Ile Met Ala Tyr Ala Gln Lys Ile Phe Lys Ile 805 810 815	2448
ttg gat gag gag aga gac aag gtg ctc acg cac att gat cac ata ttt Leu Asp Glu Glu Arg Asp Lys Val Leu Thr His Ile Asp His Ile Phe 820 825 830	2496
atg gat atc ctc act aca tgt gtg gaa aca atg tgt aat gag tac aag Met Asp Ile Leu Thr Thr Cys Val Glu Thr Met Cys Asn Glu Tyr Lys 835 840 845	2544
gtc act agt gac gct tgt atg atg acc atg tac ggg ggc atc tct ctc Val Thr Ser Asp Ala Cys Met Met Thr Met Tyr Gly Gly Ile Ser Leu 850 855 860	2592
tta agt gag ttc tgt cgg gtg ctg tgc tgc tat gtc tta gag gag act Leu Ser Glu Phe Cys Arg Val Leu Cys Cys Tyr Val Leu Glu Glu Thr 865 870 875 880	2640
agt gtg atg ctg gcc aag cgg cct ctg ata acc aag cct gag gtt atc	2688

Ser Val Met Leu Ala Lys Arg Pro Leu Ile Thr Lys Pro Glu Val Ile  
 885 890 895  
 agt gta atg aag cgc cgc att gag gag atc tgc atg aag gtc ttt gcc 2736  
 Ser Val Met Lys Arg Arg Ile Glu Glu Ile Cys Met Lys Val Phe Ala  
 900 905 910  
 cag tac att ctg ggg gcc gat cct ctg aga gtc tgc tct cct agt gtg 2784  
 Gln Tyr Ile Leu Gly Ala Asp Pro Leu Arg Val Cys Ser Pro Ser Val  
 915 920 925  
 gat gac cta cgg gcc gtc gcc gag gag tca gat gag gaa gag gct att 2832  
 Asp Asp Leu Arg Ala Val Ala Glu Glu Ser Asp Glu Glu Glu Ala Ile  
 930 935 940  
 gta gcc tac act ttg gcc acc gct ggt gtc agc tcc tct gat tct ctg 2880  
 Val Ala Tyr Thr Leu Ala Thr Ala Gly Val Ser Ser Ser Asp Ser Leu  
 945 950 955 960  
 gtg tca ccc cca gag tcc cct gta ccc gcg act atc cct ctg tcc tca 2928  
 Val Ser Pro Pro Glu Ser Pro Val Pro Ala Thr Ile Pro Leu Ser Ser  
 965 970 975  
 gta att gtg gct gag aac agt gat cag gaa gaa agt gag cag agt gat 2976  
 Val Ile Val Ala Glu Asn Ser Asp Gln Glu Glu Ser Glu Gln Ser Asp  
 980 985 990  
 gag gaa gag gag gag ggt gct cag gag gag cgg gag gac act gtg tct 3024  
 Glu Glu Glu Glu Gly Ala Gln Glu Glu Arg Glu Asp Thr Val Ser  
 995 1000 1005  
 gtc aag tct gag cca gtg tct gag ata gag gaa gtt gcc cca gag 3069  
 Val Lys Ser Glu Pro Val Ser Glu Ile Glu Glu Val Ala Pro Glu  
 1010 1015 1020  
 gaa gag gag gat ggt gct gag gaa ccc acc gcc tct gga ggc aag 3114  
 Glu Glu Glu Asp Gly Ala Glu Glu Pro Thr Ala Ser Gly Gly Lys  
 1025 1030 1035  
 agc acc cac cct atg gtg act aga agc aag gct gac cag taa 3156  
 Ser Thr His Pro Met Val Thr Arg Ser Lys Ala Asp Gln  
 1040 1045 1050  
  
 <210> 4  
 <211> 1051  
 <212> PRT  
 <213> Artificial  
  
 <220>  
 <223> hCMV pp65-IE1 fusion  
  
 <400> 4  
  
 Met Glu Ser Arg Gly Arg Arg Cys Pro Glu Met Ile Ser Val Leu Gly  
 1 5 10 15

Pro Ile Ser Gly His Val Leu Lys Ala Val Phe Ser Arg Gly Asp Thr  
 20 25 30  
 Pro Val Leu Pro His Glu Thr Arg Leu Leu Gln Thr Gly Ile His Val  
 35 40 45  
 Arg Val Ser Gln Pro Ser Leu Ile Leu Val Ser Gln Tyr Thr Pro Asp  
 50 55 60  
 Ser Thr Pro Cys His Arg Gly Asp Asn Gln Leu Gln Val Gln His Thr  
 65 70 75 80  
 Tyr Phe Thr Gly Ser Glu Val Glu Asn Val Ser Val Asn Val His Asn  
 85 90 95  
 Pro Thr Gly Arg Ser Ile Cys Pro Ser Gln Glu Pro Met Ser Ile Tyr  
 100 105 110  
 Val Tyr Ala Leu Pro Leu Lys Met Leu Asn Ile Pro Ser Ile Asn Val  
 115 120 125  
 His His Tyr Pro Ser Ala Ala Glu Arg Lys His Arg His Leu Pro Val  
 130 135 140  
 Ala Asp Ala Val Ile His Ala Ser Gly Lys Gln Met Trp Gln Ala Arg  
 145 150 155 160  
 Leu Thr Val Ser Gly Leu Ala Trp Thr Arg Gln Gln Asn Gln Trp Lys  
 165 170 175  
 Glu Pro Asp Val Tyr Tyr Thr Ser Ala Phe Val Phe Pro Thr Lys Asp  
 180 185 190  
 Val Ala Leu Arg His Val Val Cys Ala His Glu Leu Val Cys Ser Met  
 195 200 205  
 Glu Asn Thr Arg Ala Thr Lys Met Gln Val Ile Gly Asp Gln Tyr Val  
 210 215 220  
 Lys Val Tyr Leu Glu Ser Phe Cys Glu Asp Val Pro Ser Gly Lys Leu  
 225 230 235 240

Phe Met His Val Thr Leu Gly Ser Asp Val Glu Glu Asp Leu Thr Met  
 245 250 255  
 Thr Arg Asn Pro Gln Pro Phe Met Arg Pro His Glu Arg Asn Gly Phe  
 260 265 270  
 Thr Val Leu Cys Pro Lys Asn Met Ile Ile Lys Pro Gly Lys Ile Ser  
 275 280 285  
 His Ile Met Leu Asp Val Ala Phe Thr Ser His Glu His Phe Gly Leu  
 290 295 300  
 Leu Cys Pro Lys Ser Ile Pro Gly Leu Ser Ile Ser Gly Asn Leu Leu  
 305 310 315 320  
 Met Asn Gly Gln Gln Ile Phe Leu Glu Val Gln Ala Ile Arg Glu Thr  
 325 330 335  
 Val Glu Leu Arg Gln Tyr Asp Pro Val Ala Ala Leu Phe Phe Phe Asp  
 340 345 350  
 Ile Asp Leu Leu Leu Gln Arg Gly Pro Gln Tyr Ser Glu His Pro Thr  
 355 360 365  
 Phe Thr Ser Gln Tyr Arg Ile Gln Gly Lys Leu Glu Tyr Arg His Thr  
 370 375 380  
 Trp Asp Arg His Asp Glu Gly Ala Ala Gln Gly Asp Asp Asp Val Trp  
 385 390 395 400  
 Thr Ser Gly Ser Asp Ser Asp Glu Glu Leu Val Thr Thr Glu Arg Lys  
 405 410 415  
 Thr Pro Arg Val Thr Gly Gly Gly Ala Met Ala Gly Ala Ser Thr Ser  
 420 425 430  
 Ala Gly Arg Lys Arg Lys Ser Ala Ser Ser Ala Thr Ala Cys Thr Ser  
 435 440 445  
 Gly Val Met Thr Arg Gly Arg Leu Lys Ala Glu Ser Thr Val Ala Pro  
 450 455 460  
 Glu Glu Asp Thr Asp Glu Asp Ser Asp Asn Glu Ile His Asn Pro Ala



465		470		475		480
Val Phe Thr Trp	Pro Pro Trp	Gln Ala Gly	Ile Leu Ala Arg	Asn Leu		
	485		490		495	
Val Pro Met Val	Ala Thr Val	Gln Gly Gln	Asn Leu Lys	Tyr Gln Glu		
	500		505		510	
Phe Phe Trp Asp	Ala Asn Asp	Ile Tyr Arg	Ile Phe Ala	Glu Leu Glu		
	515		520		525	
Gly Val Trp Gln	Pro Ala Ala	Gln Pro Lys	Arg Arg Arg	His Arg Gln		
	530		535		540	
Asp Ala Leu Pro	Gly Pro Cys	Ile Ala Ser	Thr Pro Lys	Lys His Arg		
	545		550		555	
Gly Glu Ser Ser	Ala Lys Arg	Lys Met Asp	Pro Asp Asn	Pro Asp Glu		
	565		570		575	
Gly Pro Ser Ser	Lys Val Pro	Arg Pro Glu	Thr Pro Val	Thr Lys Ala		
	580		585		590	
Thr Thr Phe Leu	Gln Thr Met	Leu Arg Lys	Glu Val Asn	Ser Gln Leu		
	595		600		605	
Ser Leu Gly Asp	Pro Leu Phe	Pro Glu Leu	Ala Glu Glu	Ser Leu		
	610		615		620	
Thr Phe Glu Gln	Val Thr Glu	Asp Cys Asn	Glu Asn	Glu Lys Asp		
	625		630		635	
Val Leu Ala Glu	Leu Val Lys	Gln Ile Lys	Val Arg Val	Asp Met Val		
	645		650		655	
Arg His Arg Ile	Lys Glu His	Met Leu Lys	Lys Tyr Thr	Gln Thr Glu		
	660		665		670	
Glu Lys Phe Thr	Gly Ala Phe	Asn Met Met	Gly Gly Cys	Leu Gln Asn		
	675		680		685	
Ala Leu Asp Ile	Leu Asp Lys	Val His Glu	Pro Phe Glu	Glu Met Lys		
	690		695		700	

Cys Ile Gly Leu Thr Met Gln Ser Met Tyr Glu Asn Tyr Ile Val Pro  
 705 710 715 720  
 Glu Asp Lys Arg Glu Met Trp Met Ala Cys Ile Lys Glu Leu His Asp  
 725 730 735  
 Val Ser Lys Gly Ala Ala Asn Lys Leu Gly Gly Ala Leu Gln Ala Lys  
 740 745 750  
 Ala Arg Ala Lys Lys Asp Glu Leu Arg Arg Lys Met Met Tyr Met Cys  
 755 760 765  
 Tyr Arg Asn Ile Glu Phe Phe Thr Lys Asn Ser Ala Phe Pro Lys Thr  
 770 775 780  
 Thr Asn Gly Cys Ser Gln Ala Met Ala Ala Leu Gln Asn Leu Pro Gln  
 785 790 795 800  
 Cys Ser Pro Asp Glu Ile Met Ala Tyr Ala Gln Lys Ile Phe Lys Ile  
 805 810 815  
 Leu Asp Glu Glu Arg Asp Lys Val Leu Thr His Ile Asp His Ile Phe  
 820 825 830  
 Met Asp Ile Leu Thr Thr Cys Val Glu Thr Met Cys Asn Glu Tyr Lys  
 835 840 845  
 Val Thr Ser Asp Ala Cys Met Met Thr Met Tyr Gly Gly Ile Ser Leu  
 850 855 860  
 Leu Ser Glu Phe Cys Arg Val Leu Cys Cys Tyr Val Leu Glu Glu Thr  
 865 870 875 880  
 Ser Val Met Leu Ala Lys Arg Pro Leu Ile Thr Lys Pro Glu Val Ile  
 885 890 895  
 Ser Val Met Lys Arg Arg Ile Glu Glu Ile Cys Met Lys Val Phe Ala  
 900 905 910  
 Gln Tyr Ile Leu Gly Ala Asp Pro Leu Arg Val Cys Ser Pro Ser Val  
 915 920 925

Asp Asp Leu Arg Ala Val Ala Glu Glu Ser Asp Glu Glu Glu Ala Ile  
 930 935 940

Val Ala Tyr Thr Leu Ala Thr Ala Gly Val Ser Ser Ser Asp Ser Leu  
 945 950 955 960

Val Ser Pro Pro Glu Ser Pro Val Pro Ala Thr Ile Pro Leu Ser Ser  
 965 970 975

Val Ile Val Ala Glu Asn Ser Asp Gln Glu Glu Ser Glu Gln Ser Asp  
 980 985 990

Glu Glu Glu Glu Glu Gly Ala Gln Glu Glu Arg Glu Asp Thr Val Ser  
 995 1000 1005

Val Lys Ser Glu Pro Val Ser Glu Ile Glu Glu Val Ala Pro Glu  
 1010 1015 1020

Glu Glu Glu Asp Gly Ala Glu Glu Pro Thr Ala Ser Gly Gly Lys  
 1025 1030 1035

Ser Thr His Pro Met Val Thr Arg Ser Lys Ala Asp Gln  
 1040 1045 1050

<210> 5  
 <211> 15  
 <212> PRT  
 <213> Human cytomegalovirus

<400> 5

Met Glu Ser Arg Gly Arg Arg Cys Pro Glu Met Ile Ser Val Leu  
 1 5 10 15

<210> 6  
 <211> 15  
 <212> PRT  
 <213> Human cytomegalovirus

<400> 6

Gly Arg Arg Cys Pro Glu Met Ile Ser Val Leu Gly Pro Ile Ser  
 1 5 10 15

<210> 7  
 <211> 15

<212> PRT  
<213> Human cytomegalovirus

<400> 7

Pro Glu Met Ile Ser Val Leu Gly Pro Ile Ser Gly His Val Leu  
1 5 10 15

<210> 8  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 8

Ser Val Leu Gly Pro Ile Ser Gly His Val Leu Lys Ala Val Phe  
1 5 10 15

<210> 9  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 9

Pro Ile Ser Gly His Val Leu Lys Ala Val Phe Ser Arg Gly Asp  
1 5 10 15

<210> 10  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 10

His Val Leu Lys Ala Val Phe Ser Arg Gly Asp Thr Pro Val Leu  
1 5 10 15

<210> 11  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 11

Ala Val Phe Ser Arg Gly Asp Thr Pro Val Leu Pro His Glu Thr  
1 5 10 15

<210> 12  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

&lt;400&gt; 12

Arg Gly Asp Thr Pro Val Leu Pro His Glu Thr Arg Leu Leu Gln  
1 5 10 15

&lt;210&gt; 13

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 13

Pro Val Leu Pro His Glu Thr Arg Leu Leu Gln Thr Gly Ile His  
1 5 10 15

&lt;210&gt; 14

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 14

His Glu Thr Arg Leu Leu Gln Thr Gly Ile His Val Arg Val Ser  
1 5 10 15

&lt;210&gt; 15

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 15

Leu Leu Gln Thr Gly Ile His Val Arg Val Ser Gln Pro Ser Leu  
1 5 10 15

&lt;210&gt; 16

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 16

Gly Ile His Val Arg Val Ser Gln Pro Ser Leu Ile Leu Val Ser  
1 5 10 15

&lt;210&gt; 17

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 17

Arg Val Ser Gln Pro Ser Leu Ile Leu Val Ser Gln Tyr Thr Pro  
 1 5 10 15

<210> 18  
 <211> 15  
 <212> PRT  
 <213> Human cytomegalovirus

<400> 18

Pro Ser Leu Ile Leu Val Ser Gln Tyr Thr Pro Asp Ser Thr Pro  
 1 5 10 15

<210> 19  
 <211> 15  
 <212> PRT  
 <213> Human cytomegalovirus

<400> 19

Leu Val Ser Gln Tyr Thr Pro Asp Ser Thr Pro Cys His Arg Gly  
 1 5 10 15

<210> 20  
 <211> 15  
 <212> PRT  
 <213> Human cytomegalovirus

<400> 20

Tyr Thr Pro Asp Ser Thr Pro Cys His Arg Gly Asp Asn Gln Leu  
 1 5 10 15

<210> 21  
 <211> 15  
 <212> PRT  
 <213> Human cytomegalovirus

<400> 21

Ser Thr Pro Cys His Arg Gly Asp Asn Gln Leu Gln Val Gln His  
 1 5 10 15

<210> 22  
 <211> 15  
 <212> PRT  
 <213> Human cytomegalovirus

<400> 22

His Arg Gly Asp Asn Gln Leu Gln Val Gln His Thr Tyr Phe Thr

1 5 10 15

<210> 23  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 23

Asn Gln Leu Gln Val Gln His Thr Tyr Phe Thr Gly Ser Glu Val  
1 5 10 15

<210> 24  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 24

Val Gln His Thr Tyr Phe Thr Gly Ser Glu Val Glu Asn Val Ser  
1 5 10 15

<210> 25  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 25

Tyr Phe Thr Gly Ser Glu Val Glu Asn Val Ser Val Asn Val His  
1 5 10 15

<210> 26  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 26

Ser Glu Val Glu Asn Val Ser Val Asn Val His Asn Pro Thr Gly  
1 5 10 15

<210> 27  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 27

Asn Val Ser Val Asn Val His Asn Pro Thr Gly Arg Ser Ile Cys  
1 5 10 15

<210> 28  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 28

Asn Val His Asn Pro Thr Gly Arg Ser Ile Cys Pro Ser Gln Glu  
1 5 10 15

<210> 29  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 29

Pro Thr Gly Arg Ser Ile Cys Pro Ser Gln Glu Pro Met Ser Ile  
1 5 10 15

<210> 30  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 30

Ser Ile Cys Pro Ser Gln Glu Pro Met Ser Ile Tyr Val Tyr Ala  
1 5 10 15

<210> 31  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 31

Ser Gln Glu Pro Met Ser Ile Tyr Val Tyr Ala Leu Pro Leu Lys  
1 5 10 15

<210> 32  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 32

Met Ser Ile Tyr Val Tyr Ala Leu Pro Leu Lys Met Leu Asn Ile  
1 5 10 15

<210> 33



<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 33

Val Tyr Ala Leu Pro Leu Lys Met Leu Asn Ile Pro Ser Ile Asn  
1 5 10 15

<210> 34  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 34

Pro Leu Lys Met Leu Asn Ile Pro Ser Ile Asn Val His His Tyr  
1 5 10 15

<210> 35  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 35

Leu Asn Ile Pro Ser Ile Asn Val His His Tyr Pro Ser Ala Ala  
1 5 10 15

<210> 36  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 36

Ser Ile Asn Val His His Tyr Pro Ser Ala Ala Glu Arg Lys His  
1 5 10 15

<210> 37  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 37

His His Tyr Pro Ser Ala Ala Glu Arg Lys His Arg His Leu Pro  
1 5 10 15

<210> 38  
<211> 15  
<212> PRT

<213> Human cytomegalovirus

<400> 38

Ser Ala Ala Glu Arg Lys His Arg His Leu Pro Val Ala Asp Ala  
1 5 10 15

<210> 39

<211> 15

<212> PRT

<213> Human cytomegalovirus

<400> 39

Arg Lys His Arg His Leu Pro Val Ala Asp Ala Val Ile His Ala  
1 5 10 15

<210> 40

<211> 15

<212> PRT

<213> Human cytomegalovirus

<400> 40

His Leu Pro Val Ala Asp Ala Val Ile His Ala Ser Gly Lys Gln  
1 5 10 15

<210> 41

<211> 15

<212> PRT

<213> Human cytomegalovirus

<400> 41

Ala Asp Ala Val Ile His Ala Ser Gly Lys Gln Met Trp Gln Ala  
1 5 10 15

<210> 42

<211> 15

<212> PRT

<213> Human cytomegalovirus

<400> 42

Ile His Ala Ser Gly Lys Gln Met Trp Gln Ala Arg Leu Thr Val  
1 5 10 15

<210> 43

<211> 15

<212> PRT

<213> Human cytomegalovirus

&lt;400&gt; 43

Gly Lys Gln Met Trp Gln Ala Arg Leu Thr Val Ser Gly Leu Ala  
1 5 10 15

&lt;210&gt; 44

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 44

Trp Gln Ala Arg Leu Thr Val Ser Gly Leu Ala Trp Thr Arg Gln  
1 5 10 15

&lt;210&gt; 45

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 45

Leu Thr Val Ser Gly Leu Ala Trp Thr Arg Gln Gln Asn Gln Trp  
1 5 10 15

&lt;210&gt; 46

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 46

Gly Leu Ala Trp Thr Arg Gln Gln Asn Gln Trp Lys Glu Pro Asp  
1 5 10 15

&lt;210&gt; 47

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 47

Thr Arg Gln Gln Asn Gln Trp Lys Glu Pro Asp Val Tyr Tyr Thr  
1 5 10 15

&lt;210&gt; 48

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 48

Asn Gln Trp Lys Glu Pro Asp Val Tyr Tyr Thr Ser Ala Phe Val  
 1 5 10 15

<210> 49  
 <211> 15  
 <212> PRT  
 <213> Human cytomegalovirus  
 <400> 49

Glu Pro Asp Val Tyr Tyr Thr Ser Ala Phe Val Phe Pro Thr Lys  
 1 5 10 15

<210> 50  
 <211> 15  
 <212> PRT  
 <213> Human cytomegalovirus  
 <400> 50

Tyr Tyr Thr Ser Ala Phe Val Phe Pro Thr Lys Asp Val Ala Leu  
 1 5 10 15

<210> 51  
 <211> 15  
 <212> PRT  
 <213> Human cytomegalovirus  
 <400> 51

Ala Phe Val Phe Pro Thr Lys Asp Val Ala Leu Arg His Val Val  
 1 5 10 15

<210> 52  
 <211> 15  
 <212> PRT  
 <213> Human cytomegalovirus  
 <400> 52

Pro Thr Lys Asp Val Ala Leu Arg His Val Val Cys Ala His Glu  
 1 5 10 15

<210> 53  
 <211> 15  
 <212> PRT  
 <213> Human cytomegalovirus  
 <400> 53

Val Ala Leu Arg His Val Val Cys Ala His Glu Leu Val Cys Ser  
 1 5 10 15

<210> 54  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 54

His Val Val Cys Ala His Glu Leu Val Cys Ser Met Glu Asn Thr  
1 5 10 15

<210> 55  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 55

Ala His Glu Leu Val Cys Ser Met Glu Asn Thr Arg Ala Thr Lys  
1 5 10 15

<210> 56  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 56

Val Cys Ser Met Glu Asn Thr Arg Ala Thr Lys Met Gln Val Ile  
1 5 10 15

<210> 57  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 57

Glu Asn Thr Arg Ala Thr Lys Met Gln Val Ile Gly Asp Gln Tyr  
1 5 10 15

<210> 58  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 58

Ala Thr Lys Met Gln Val Ile Gly Asp Gln Tyr Val Lys Val Tyr  
1 5 10 15

<210> 59  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 59

Gln Val Ile Gly Asp Gln Tyr Val Lys Val Tyr Leu Glu Ser Phe  
1 5 10 15

<210> 60  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 60

Asp Gln Tyr Val Lys Val Tyr Leu Glu Ser Phe Cys Glu Asp Val  
1 5 10 15

<210> 61  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 61

Lys Val Tyr Leu Glu Ser Phe Cys Glu Asp Val Pro Ser Gly Lys  
1 5 10 15

<210> 62  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 62

Glu Ser Phe Cys Glu Asp Val Pro Ser Gly Lys Leu Phe Met His  
1 5 10 15

<210> 63  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 63

Glu Asp Val Pro Ser Gly Lys Leu Phe Met His Val Thr Leu Gly  
1 5 10 15

<210> 64  
<211> 15

<212> PRT  
<213> Human cytomegalovirus

<400> 64

Ser Gly Lys Leu Phe Met His Val Thr Leu Gly Ser Asp Val Glu  
1 5 10 15

<210> 65  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 65

Phe Met His Val Thr Leu Gly Ser Asp Val Glu Glu Asp Leu Thr  
1 5 10 15

<210> 66  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 66

Thr Leu Gly Ser Asp Val Glu Glu Asp Leu Thr Met Thr Arg Asn  
1 5 10 15

<210> 67  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 67

Asp Val Glu Glu Asp Leu Thr Met Thr Arg Asn Pro Gln Pro Phe  
1 5 10 15

<210> 68  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 68

Asp Leu Thr Met Thr Arg Asn Pro Gln Pro Phe Met Arg Pro His  
1 5 10 15

<210> 69  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

&lt;400&gt; 69

Thr	Arg	Asn	Pro	Gln	Pro	Phe	Met	Arg	Pro	His	Glu	Arg	Asn	Gly
1				5					10					15

&lt;210&gt; 70

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 70

Gln	Pro	Phe	Met	Arg	Pro	His	Glu	Arg	Asn	Gly	Phe	Thr	Val	Leu
1				5					10					15

&lt;210&gt; 71

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 71

Arg	Pro	His	Glu	Arg	Asn	Gly	Phe	Thr	Val	Leu	Cys	Pro	Lys	Asn
1				5					10					15

&lt;210&gt; 72

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 72

Arg	Asn	Gly	Phe	Thr	Val	Leu	Cys	Pro	Lys	Asn	Met	Ile	Ile	Lys
1				5					10					15

&lt;210&gt; 73

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 73

Thr	Val	Leu	Cys	Pro	Lys	Asn	Met	Ile	Ile	Lys	Pro	Gly	Lys	Ile
1				5					10					15

&lt;210&gt; 74

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 74



Pro Lys Asn Met Ile Ile Lys Pro Gly Lys Ile Ser His Ile Met  
1 5 10 15

<210> 75  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 75

Ile Ile Lys Pro Gly Lys Ile Ser His Ile Met Leu Asp Val Ala  
1 5 10 15

<210> 76  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 76

Gly Lys Ile Ser His Ile Met Leu Asp Val Ala Phe Thr Ser His  
1 5 10 15

<210> 77  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 77

His Ile Met Leu Asp Val Ala Phe Thr Ser His Glu His Phe Gly  
1 5 10 15

<210> 78  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 78

Asp Val Ala Phe Thr Ser His Glu His Phe Gly Leu Leu Cys Pro  
1 5 10 15

<210> 79  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 79

Thr Ser His Glu His Phe Gly Leu Leu Cys Pro Lys Ser Ile Pro

1 5 10 15

<210> 80  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 80

His Phe Gly Leu Leu Cys Pro Lys Ser Ile Pro Gly Leu Ser Ile  
1 5 10 15

<210> 81  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 81

Leu Cys Pro Lys Ser Ile Pro Gly Leu Ser Ile Ser Gly Asn Leu  
1 5 10 15

<210> 82  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 82

Ser Ile Pro Gly Leu Ser Ile Ser Gly Asn Leu Leu Met Asn Gly  
1 5 10 15

<210> 83  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 83

Leu Ser Ile Ser Gly Asn Leu Leu Met Asn Gly Gln Gln Ile Phe  
1 5 10 15

<210> 84  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 84

Gly Asn Leu Leu Met Asn Gly Gln Gln Ile Phe Leu Glu Val Gln  
1 5 10 15

<210> 85  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 85

Met Asn Gly Gln Gln Ile Phe Leu Glu Val Gln Ala Ile Arg Glu  
1 5 10 15

<210> 86  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 86

Gln Ile Phe Leu Glu Val Gln Ala Ile Arg Glu Thr Val Glu Leu  
1 5 10 15

<210> 87  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 87

Glu Val Gln Ala Ile Arg Glu Thr Val Glu Leu Arg Gln Tyr Asp  
1 5 10 15

<210> 88  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 88

Ile Arg Glu Thr Val Glu Leu Arg Gln Tyr Asp Pro Val Ala Ala  
1 5 10 15

<210> 89  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 89

Val Glu Leu Arg Gln Tyr Asp Pro Val Ala Ala Leu Phe Phe Phe  
1 5 10 15

<210> 90

<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 90

Gln Tyr Asp Pro Val Ala Ala Leu Phe Phe Phe Asp Ile Asp Leu  
1 5 10 15

<210> 91  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 91

Val Ala Ala Leu Phe Phe Phe Asp Ile Asp Leu Leu Leu Gln Arg  
1 5 10 15

<210> 92  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 92

Phe Phe Phe Asp Ile Asp Leu Leu Leu Gln Arg Gly Pro Gln Tyr  
1 5 10 15

<210> 93  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 93

Ile Asp Leu Leu Leu Gln Arg Gly Pro Gln Tyr Ser Glu His Pro  
1 5 10 15

<210> 94  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 94

Leu Gln Arg Gly Pro Gln Tyr Ser Glu His Pro Thr Phe Thr Ser  
1 5 10 15

<210> 95  
<211> 15  
<212> PRT

<213> Human cytomegalovirus

<400> 95

Pro Gln Tyr Ser Glu His Pro Thr Phe Thr Ser Gln Tyr Arg Ile  
1 5 10 15

<210> 96

<211> 15

<212> PRT

<213> Human cytomegalovirus

<400> 96

Glu His Pro Thr Phe Thr Ser Gln Tyr Arg Ile Gln Gly Lys Leu  
1 5 10 15

<210> 97

<211> 15

<212> PRT

<213> Human cytomegalovirus

<400> 97

Phe Thr Ser Gln Tyr Arg Ile Gln Gly Lys Leu Glu Tyr Arg His  
1 5 10 15

<210> 98

<211> 15

<212> PRT

<213> Human cytomegalovirus

<400> 98

Tyr Arg Ile Gln Gly Lys Leu Glu Tyr Arg His Thr Trp Asp Arg  
1 5 10 15

<210> 99

<211> 15

<212> PRT

<213> Human cytomegalovirus

<400> 99

Gly Lys Leu Glu Tyr Arg His Thr Trp Asp Arg His Asp Glu Gly  
1 5 10 15

<210> 100

<211> 15

<212> PRT

<213> Human cytomegalovirus

&lt;400&gt; 100

Tyr Arg His Thr Trp Asp Arg His Asp Glu Gly Ala Ala Gln Gly  
1 5 10 15

&lt;210&gt; 101

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 101

Trp Asp Arg His Asp Glu Gly Ala Ala Gln Gly Asp Asp Asp Val  
1 5 10 15

&lt;210&gt; 102

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 102

Asp Glu Gly Ala Ala Gln Gly Asp Asp Asp Val Trp Thr Ser Gly  
1 5 10 15

&lt;210&gt; 103

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 103

Ala Gln Gly Asp Asp Asp Val Trp Thr Ser Gly Ser Asp Ser Asp  
1 5 10 15

&lt;210&gt; 104

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 104

Asp Asp Val Trp Thr Ser Gly Ser Asp Ser Asp Glu Glu Leu Val  
1 5 10 15

&lt;210&gt; 105

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 105

Thr Ser Gly Ser Asp Ser Asp Glu Glu Leu Val Thr Thr Glu Arg  
1 5 10 15

<210> 106  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 106

Asp Ser Asp Glu Glu Leu Val Thr Thr Glu Arg Lys Thr Pro Arg  
1 5 10 15

<210> 107  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 107

Glu Leu Val Thr Thr Glu Arg Lys Thr Pro Arg Val Thr Gly Gly  
1 5 10 15

<210> 108  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 108

Thr Glu Arg Lys Thr Pro Arg Val Thr Gly Gly Gly Ala Met Ala  
1 5 10 15

<210> 109  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 109

Thr Pro Arg Val Thr Gly Gly Gly Ala Met Ala Gly Ala Ser Thr  
1 5 10 15

<210> 110  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 110

Thr Gly Gly Gly Ala Met Ala Gly Ala Ser Thr Ser Ala Gly Arg  
1 5 10 15

<210> 111  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 111

Ala Met Ala Gly Ala Ser Thr Ser Ala Gly Arg Lys Arg Lys Ser  
1                   5                   10                   15

<210> 112  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 112

Ala Ser Thr Ser Ala Gly Arg Lys Arg Lys Ser Ala Ser Ser Ala  
1                   5                   10                   15

<210> 113  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 113

Ala Gly Arg Lys Arg Lys Ser Ala Ser Ser Ala Thr Ala Cys Thr  
1                   5                   10                   15

<210> 114  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 114

Arg Lys Ser Ala Ser Ser Ala Thr Ala Cys Thr Ser Gly Val Met  
1                   5                   10                   15

<210> 115  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 115

Ser Ser Ala Thr Ala Cys Thr Ser Gly Val Met Thr Arg Gly Arg  
1                   5                   10                   15



<210> 116  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 116

Ala Cys Thr Ser Gly Val Met Thr Arg Gly Arg Leu Lys Ala Glu  
1 5 10 15

<210> 117  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 117

Gly Val Met Thr Arg Gly Arg Leu Lys Ala Glu Ser Thr Val Ala  
1 5 10 15

<210> 118  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 118

Arg Gly Arg Leu Lys Ala Glu Ser Thr-Val Ala Pro Glu Glu Asp  
1 5 10 15

<210> 119  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 119

Lys Ala Glu Ser Thr Val Ala Pro Glu Glu Asp Thr Asp Glu Asp  
1 5 10 15

<210> 120  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 120

Thr Val Ala Pro Glu Glu Asp Thr Asp Glu Asp Ser Asp Asn Glu  
1 5 10 15

<210> 121  
<211> 15

<212> PRT  
<213> Human cytomegalovirus

<400> 121

Glu	Glu	Asp	Thr	Asp	Glu	Asp	Ser	Asp	Asn	Glu	Ile	His	Asn	Pro
1				5					10					15

<210> 122  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 122

Asp	Glu	Asp	Ser	Asp	Asn	Glu	Ile	His	Asn	Pro	Ala	Val	Phe	Thr
1				5					10					15

<210> 123  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 123

Asp	Asn	Glu	Ile	His	Asn	Pro	Ala	Val	Phe	Thr	Trp	Pro	Pro	Trp
1				5					10					15

<210> 124  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 124

His	Asn	Pro	Ala	Val	Phe	Thr	Trp	Pro	Pro	Trp	Gln	Ala	Gly	Ile
1				5					10					15

<210> 125  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 125

Val	Phe	Thr	Trp	Pro	Pro	Trp	Gln	Ala	Gly	Ile	Leu	Ala	Arg	Asn
1				5					10					15

<210> 126  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

&lt;400&gt; 126

Pro Pro Trp Gln Ala Gly Ile Leu Ala Arg Asn Leu Val Pro Met  
1 5 10 15

&lt;210&gt; 127

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 127

Ala Gly Ile Leu Ala Arg Asn Leu Val Pro Met Val Ala Thr Val  
1 5 10 15

&lt;210&gt; 128

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 128

Ala Arg Asn Leu Val Pro Met Val Ala Thr Val Gln Gly Gln Asn  
1 5 10 15

&lt;210&gt; 129

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 129

Val Pro Met Val Ala Thr Val Gln Gly Gln Asn Leu Lys Tyr Gln  
1 5 10 15

&lt;210&gt; 130

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 130

Ala Thr Val Gln Gly Gln Asn Leu Lys Tyr Gln Glu Phe Phe Trp  
1 5 10 15

&lt;210&gt; 131

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 131

Gly Gln Asn Leu Lys Tyr Gln Glu Phe Phe Trp Asp Ala Asn Asp  
1 5 10 15

<210> 132  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 132

Lys Tyr Gln Glu Phe Phe Trp Asp Ala Asn Asp Ile Tyr Arg Ile  
1 5 10 15

<210> 133  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 133

Phe Phe Trp Asp Ala Asn Asp Ile Tyr Arg Ile Phe Ala Glu Leu  
1 5 10 15

<210> 134  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 134

Ala Asn Asp Ile Tyr Arg Ile Phe Ala Glu Leu Glu Gly Val Trp  
1 5 10 15

<210> 135  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 135

Tyr Arg Ile Phe Ala Glu Leu Glu Gly Val Trp Gln Pro Ala Ala  
1 5 10 15

<210> 136  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 136

Ala Glu Leu Glu Gly Val Trp Gln Pro Ala Ala Gln Pro Lys Arg

1 5 10 15

<210> 137  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 137

Gly Val Trp Gln Pro Ala Ala Gln Pro Lys Arg Arg Arg His Arg  
1 5 10 15

<210> 138  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 138

Pro Ala Ala Gln Pro Lys Arg Arg Arg His Arg Gln Asp Ala Leu  
1 5 10 15

<210> 139  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 139

Pro Lys Arg Arg Arg His Arg Gln Asp Ala Leu Pro Gly Pro Cys  
1 5 10 15

<210> 140  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 140

Arg His Arg Gln Asp Ala Leu Pro Gly Pro Cys Ile Ala Ser Thr  
1 5 10 15

<210> 141  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 141

Asp Ala Leu Pro Gly Pro Cys Ile Ala Ser Thr Pro Lys Lys His  
1 5 10 15

<210> 142  
<211> 13  
<212> PRT  
<213> Human cytomegalovirus

<400> 142

Gly Pro Cys Ile Ala Ser Thr Pro Lys Lys His Arg Gly  
1 5 10

<210> 143  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 143

Met Glu Ser Ser Ala Lys Arg Lys Met Asp Pro Asp Asn Pro Asp  
1 5 10 15

<210> 144  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 144

Ala Lys Arg Lys Met Asp Pro Asp Asn Pro Asp Glu Gly Pro Ser  
1 5 10 15

<210> 145  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 145

Met Asp Pro Asp Asn Pro Asp Glu Gly Pro Ser Ser Lys Val Pro  
1 5 10 15

<210> 146  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 146

Asn Pro Asp Glu Gly Pro Ser Ser Lys Val Pro Arg Pro Glu Thr  
1 5 10 15

<210> 147

<211> 15  
 <212> PRT  
 <213> Human cytomegalovirus

<400> 147

Gly	Pro	Ser	Ser	Lys	Val	Pro	Arg	Pro	Glu	Thr	Pro	Val	Thr	Lys
1				5					10					15

<210> 148  
 <211> 15  
 <212> PRT  
 <213> Human cytomegalovirus

<400> 148

Lys	Val	Pro	Arg	Pro	Glu	Thr	Pro	Val	Thr	Lys	Ala	Thr	Thr	Phe
1				5					10					15

<210> 149  
 <211> 15  
 <212> PRT  
 <213> Human cytomegalovirus

<400> 149

Pro	Glu	Thr	Pro	Val	Thr	Lys	Ala	Thr	Thr	Phe	Leu	Gln	Thr	Met
1				5					10					15

<210> 150  
 <211> 15  
 <212> PRT  
 <213> Human cytomegalovirus

<400> 150

Val	Thr	Lys	Ala	Thr	Thr	Phe	Leu	Gln	Thr	Met	Leu	Arg	Lys	Glu
1				5					10					15

<210> 151  
 <211> 15  
 <212> PRT  
 <213> Human cytomegalovirus

<400> 151

Thr	Thr	Phe	Leu	Gln	Thr	Met	Leu	Arg	Lys	Glu	Val	Asn	Ser	Gln
1				5					10					15

<210> 152  
 <211> 15  
 <212> PRT

<213> Human cytomegalovirus

<400> 152

Gln Thr Met Leu Arg Lys Glu Val Asn Ser Gln Leu Ser Leu Gly  
1 5 10 15

<210> 153

<211> 15

<212> PRT

<213> Human cytomegalovirus

<400> 153

Arg Lys Glu Val Asn Ser Gln Leu Ser Leu Gly Asp Pro Leu Phe  
1 5 10 15

<210> 154

<211> 15

<212> PRT

<213> Human cytomegalovirus

<400> 154

Asn Ser Gln Leu Ser Leu Gly Asp Pro Leu Phe Pro Glu Leu Ala  
1 5 10 15

<210> 155

<211> 15

<212> PRT

<213> Human cytomegalovirus

<400> 155

Ser Leu Gly Asp Pro Leu Phe Pro Glu Leu Ala Glu Glu Ser Leu  
1 5 10 15

<210> 156

<211> 15

<212> PRT

<213> Human cytomegalovirus

<400> 156

Pro Leu Phe Pro Glu Leu Ala Glu Glu Ser Leu Lys Thr Phe Glu  
1 5 10 15

<210> 157

<211> 15

<212> PRT

<213> Human cytomegalovirus



&lt;400&gt; 157

Glu Leu Ala Glu Glu Ser Leu Lys Thr Phe Glu Gln Val Thr Glu  
1 5 10 15

&lt;210&gt; 158

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 158

Glu Ser Leu Lys Thr Phe Glu Gln Val Thr Glu Asp Cys Asn Glu  
1 5 10 15

&lt;210&gt; 159

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 159

Thr Phe Glu Gln Val Thr Glu Asp Cys Asn Glu Asn Pro Glu Lys  
1 5 10 15

&lt;210&gt; 160

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 160

Val Thr Glu Asp Cys Asn Glu Asn Pro Glu Lys Asp Val Leu Ala  
1 5 10 15

&lt;210&gt; 161

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 161

Cys Asn Glu Asn Pro Glu Lys Asp Val Leu Ala Glu Leu Val Lys  
1 5 10 15

&lt;210&gt; 162

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 162

Pro Glu Lys Asp Val Leu Ala Glu Leu Val Lys Gln Ile Lys Val  
1 5 10 15

<210> 163  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 163

Val Leu Ala Glu Leu Val Lys Gln Ile Lys Val Arg Val Asp Met  
1 5 10 15

<210> 164  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 164

Leu Val Lys Gln Ile Lys Val Arg Val Asp Met Val Arg His Arg  
1 5 10 15

<210> 165  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 165

Ile Lys Val Arg Val Asp Met Val Arg His Arg Ile Lys Glu His  
1 5 10 15

<210> 166  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 166

Val Asp Met Val Arg His Arg Ile Lys Glu His Met Leu Lys Lys  
1 5 10 15

<210> 167  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 167

Arg His Arg Ile Lys Glu His Met Leu Lys Lys Tyr Thr Gln Thr  
1 5 10 15

<210> 168  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 168

Lys	Glu	His	Met	Leu	Lys	Lys	Tyr	Thr	Gln	Thr	Glu	Glu	Lys	Phe
1				5					10					15

<210> 169  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 169

Leu	Lys	Lys	Tyr	Thr	Gln	Thr	Glu	Glu	Lys	Phe	Thr	Gly	Ala	Phe
1				5					10					15

<210> 170  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 170

Thr	Gln	Thr	Glu	Glu	Lys	Phe	Thr	Gly	Ala	Phe	Asn	Met	Met	Gly
1				5					10					15

<210> 171  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 171

Glu	Lys	Phe	Thr	Gly	Ala	Phe	Asn	Met	Met	Gly	Gly	Cys	Leu	Gln
1				5					10					15

<210> 172  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 172

Gly	Ala	Phe	Asn	Met	Met	Gly	Gly	Cys	Leu	Gln	Asn	Ala	Leu	Asp
1				5					10					15

<210> 173  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 173

Met Met Gly Gly Cys Leu Gln Asn Ala Leu Asp Ile Leu Asp Lys  
1 5 10 15

<210> 174  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 174

Cys Leu Gln Asn Ala Leu Asp Ile Leu Asp Lys Val His Glu Pro  
1 5 10 15

<210> 175  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 175

Ala Leu Asp Ile Leu Asp Lys Val His Glu Pro Phe Glu Glu Met  
1 5 10 15

<210> 176  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 176

Leu Asp Lys Val His Glu Pro Phe Glu Glu Met Lys Cys Ile Gly  
1 5 10 15

<210> 177  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 177

His Glu Pro Phe Glu Glu Met Lys Cys Ile Gly Leu Thr Met Gln  
1 5 10 15

<210> 178  
<211> 15

<212> PRT  
<213> Human cytomegalovirus

<400> 178

Glu Glu Met Lys Cys Ile Gly Leu Thr Met Gln Ser Met Tyr Glu  
1 5 10 15

<210> 179  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 179

Cys Ile Gly Leu Thr Met Gln Ser Met Tyr Glu Asn Tyr Ile Val  
1 5 10 15

<210> 180  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 180

Thr Met Gln Ser Met Tyr Glu Asn Tyr Ile Val Pro Glu Asp Lys  
1 5 10 15

<210> 181  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 181

Met Tyr Glu Asn Tyr Ile Val Pro Glu Asp Lys Arg Glu Met Trp  
1 5 10 15

<210> 182  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 182

Tyr Ile Val Pro Glu Asp Lys Arg Glu Met Trp Met Ala Cys Ile  
1 5 10 15

<210> 183  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

&lt;400&gt; 183

Glu Asp Lys Arg Glu Met Trp Met Ala Cys Ile Lys Glu Leu His  
1 5 10 15

&lt;210&gt; 184

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 184

Glu Met Trp Met Ala Cys Ile Lys Glu Leu His Asp Val Ser Lys  
1 5 10 15

&lt;210&gt; 185

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 185

Ala Cys Ile Lys Glu Leu His Asp Val Ser Lys Gly Ala Ala Asn  
1 5 10 15

&lt;210&gt; 186

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 186

Glu Leu His Asp Val Ser Lys Gly Ala Ala Asn Lys Leu Gly Gly  
1 5 10 15

&lt;210&gt; 187

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 187

Val Ser Lys Gly Ala Ala Asn Lys Leu Gly Gly Ala Leu Gln Ala  
1 5 10 15

&lt;210&gt; 188

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 188

Ala Ala Asn Lys Leu Gly Gly Ala Leu Gln Ala Lys Ala Arg Ala  
1 5 10 15

<210> 189  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 189

Leu Gly Gly Ala Leu Gln Ala Lys Ala Arg Ala Lys Lys Asp Glu  
1 5 10 15

<210> 190  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 190

Leu Gln Ala Lys Ala Arg Ala Lys Lys Asp Glu Leu Arg Arg Lys  
1 5 10 15

<210> 191  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 191

Ala Arg Ala Lys Lys Asp Glu Leu Arg Arg Lys Met Met Tyr Met  
1 5 10 15

<210> 192  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 192

Lys Asp Glu Leu Arg Arg Lys Met Met Tyr Met Cys Tyr Arg Asn  
1 5 10 15

<210> 193  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 193

Arg Arg Lys Met Met Tyr Met Cys Tyr Arg Asn Ile Glu Phe Phe

1 5 10 15

<210> 194  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 194

Met Tyr Met Cys Tyr Arg Asn Ile Glu Phe Phe Thr Lys Asn Ser  
1 5 10 15

<210> 195  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 195

Tyr Arg Asn Ile Glu Phe Phe Thr Lys Asn Ser Ala Phe Pro Lys  
1 5 10 15

<210> 196  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 196

Glu Phe Phe Thr Lys Asn Ser Ala Phe Pro Lys Thr Thr Asn Gly  
1 5 10 15

<210> 197  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 197

Lys Asn Ser Ala Phe Pro Lys Thr Thr Asn Gly Cys Ser Gln Ala  
1 5 10 15

<210> 198  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 198

Phe Pro Lys Thr Thr Asn Gly Cys Ser Gln Ala Met Ala Ala Leu  
1 5 10 15



<210> 199  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 199

Thr Asn Gly Cys Ser Gln Ala Met Ala Ala Leu Gln Asn Leu Pro  
1 5 10 15

<210> 200  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 200

Ser Gln Ala Met Ala Ala Leu Gln Asn Leu Pro Gln Cys Ser Pro  
1 5 10 15

<210> 201  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 201

Ala Ala Leu Gln Asn Leu Pro Gln Cys Ser Pro Asp Glu Ile Met  
1 5 10 15

<210> 202  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 202

Asn Leu Pro Gln Cys Ser Pro Asp Glu Ile Met Ala Tyr Ala Gln  
1 5 10 15

<210> 203  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 203

Cys Ser Pro Asp Glu Ile Met Ala Tyr Ala Gln Lys Ile Phe Lys  
1 5 10 15

<210> 204

<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 204

Glu Ile Met Ala Tyr Ala Gln Lys Ile Phe Lys Ile Leu Asp Glu  
1 5 10 15

<210> 205  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 205

Tyr Ala Gln Lys Ile Phe Lys Ile Leu Asp Glu Glu Arg Asp Lys  
1 5 10 15

<210> 206  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 206

Ile Phe Lys Ile Leu Asp Glu Glu Arg Asp Lys Val Leu Thr His  
1 5 10 15

<210> 207  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 207

Leu Asp Glu Glu Arg Asp Lys Val Leu Thr His Ile Asp His Ile  
1 5 10 15

<210> 208  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 208

Arg Asp Lys Val Leu Thr His Ile Asp His Ile Phe Met Asp Ile  
1 5 10 15

<210> 209  
<211> 15  
<212> PRT

<213> Human cytomegalovirus

<400> 209

Leu	Thr	His	Ile	Asp	His	Ile	Phe	Met	Asp	Ile	Leu	Thr	Thr	Cys
1				5					10					15

<210> 210

<211> 15

<212> PRT

<213> Human cytomegalovirus

<400> 210

Asp	His	Ile	Phe	Met	Asp	Ile	Leu	Thr	Thr	Cys	Val	Glu	Thr	Met
1				5					10					15

<210> 211

<211> 15

<212> PRT

<213> Human cytomegalovirus

<400> 211

Met	Asp	Ile	Leu	Thr	Thr	Cys	Val	Glu	Thr	Met	Cys	Asn	Glu	Tyr
1				5					10					15

<210> 212

<211> 15

<212> PRT

<213> Human cytomegalovirus

<400> 212

Thr	Thr	Cys	Val	Glu	Thr	Met	Cys	Asn	Glu	Tyr	Lys	Val	Thr	Ser
1				5					10					15

<210> 213

<211> 15

<212> PRT

<213> Human cytomegalovirus

<400> 213

Glu	Thr	Met	Cys	Asn	Glu	Tyr	Lys	Val	Thr	Ser	Asp	Ala	Cys	Met
1				5					10					15

<210> 214

<211> 15

<212> PRT

<213> Human cytomegalovirus

&lt;400&gt; 214

Asn Glu Tyr Lys Val Thr Ser Asp Ala Cys Met Met Thr Met Tyr  
1 5 10 15

&lt;210&gt; 215

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 215

Val Thr Ser Asp Ala Cys Met Met Thr Met Tyr Gly Gly Ile Ser  
1 5 10 15

&lt;210&gt; 216

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 216

Ala Cys Met Met Thr Met Tyr Gly Gly Ile Ser Leu Leu Ser Glu  
1 5 10 15

&lt;210&gt; 217

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 217

Thr Met Tyr Gly Gly Ile Ser Leu Leu Ser Glu Phe Cys Arg Val  
1 5 10 15

&lt;210&gt; 218

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 218

Gly Ile Ser Leu Leu Ser Glu Phe Cys Arg Val Leu Cys Cys Tyr  
1 5 10 15

&lt;210&gt; 219

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 219

Leu Ser Glu Phe Cys Arg Val Leu Cys Cys Tyr Val Leu Glu Glu  
1 5 10 15

<210> 220  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 220

Cys Arg Val Leu Cys Cys Tyr Val Leu Glu Glu Thr Ser Val Met  
1 5 10 15

<210> 221  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 221

Cys Cys Tyr Val Leu Glu Glu Thr Ser Val Met Leu Ala Lys Arg  
1 5 10 15

<210> 222  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 222

Leu Glu Glu Thr Ser Val Met Leu Ala Lys Arg Pro Leu Ile Thr  
1 5 10 15

<210> 223  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 223

Ser Val Met Leu Ala Lys Arg Pro Leu Ile Thr Lys Pro Glu Val  
1 5 10 15

<210> 224  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 224

Ala Lys Arg Pro Leu Ile Thr Lys Pro Glu Val Ile Ser Val Met  
1 5 10 15

<210> 225  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 225

Leu Ile Thr Lys Pro Glu Val Ile Ser Val Met Lys Arg Arg Ile  
1 5 10 15

<210> 226  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 226

Pro Glu Val Ile Ser Val Met Lys Arg Arg Ile Glu Glu Ile Cys  
1 5 10 15

<210> 227  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 227

Ser Val Met Lys Arg Arg Ile Glu Glu Ile Cys Met Lys Val Phe  
1 5 10 15

<210> 228  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 228

Arg Arg Ile Glu Glu Ile Cys Met Lys Val Phe Ala Gln Tyr Ile  
1 5 10 15

<210> 229  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 229

Glu Ile Cys Met Lys Val Phe Ala Gln Tyr Ile Leu Gly Ala Asp  
1 5 10 15

<210> 230  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 230

Lys Val Phe Ala Gln Tyr Ile Leu Gly Ala Asp Pro Leu Arg Val  
1 5 10 15

<210> 231  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 231

Gln Tyr Ile Leu Gly Ala Asp Pro Leu Arg Val Cys Ser Pro Ser  
1 5 10 15

<210> 232  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 232

Gly Ala Asp Pro Leu Arg Val Cys Ser Pro Ser Val Asp Asp Leu  
1 5 10 15

<210> 233  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 233

Leu Arg Val Cys Ser Pro Ser Val Asp Asp Leu Arg Ala Ile Ala  
1 5 10 15

<210> 234  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 234

Ser Pro Ser Val Asp Asp Leu Arg Ala Ile Ala Glu Glu Ser Asp  
1 5 10 15

<210> 235  
<211> 15

<212> PRT  
<213> Human cytomegalovirus

<400> 235

Asp Asp Leu Arg Ala Ile Ala Glu Glu Ser Asp Glu Glu Glu Ala  
1 5 10 15

<210> 236  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 236

Ala Ile Ala Glu Glu Ser Asp Glu Glu Glu Ala Ile Val Ala Tyr  
1 5 10 15

<210> 237  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 237

Glu Ser Asp Glu Glu Glu Ala Ile Val Ala Tyr Thr Leu Ala Thr  
1 5 10 15

<210> 238  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 238

Glu Glu Ala Ile Val Ala Tyr Thr Leu Ala Thr Ala Gly Val Ser  
1 5 10 15

<210> 239  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 239

Val Ala Tyr Thr Leu Ala Thr Ala Gly Val Ser Ser Ser Asp Ser  
1 5 10 15

<210> 240  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus



&lt;400&gt; 240

Leu Ala Thr Ala Gly Val Ser Ser Ser Asp Ser Leu Val Ser Pro  
1 5 10 15

&lt;210&gt; 241

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 241

Gly Val Ser Ser Ser Asp Ser Leu Val Ser Pro Pro Glu Ser Pro  
1 5 10 15

&lt;210&gt; 242

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 242

Ser Asp Ser Leu Val Ser Pro Pro Glu Ser Pro Val Pro Ala Thr  
1 5 10 15

&lt;210&gt; 243

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 243

Val Ser Pro Pro Glu Ser Pro Val Pro Ala Thr Ile Pro Leu Ser  
1 5 10 15

&lt;210&gt; 244

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 244

Glu Ser Pro Val Pro Ala Thr Ile Pro Leu Ser Ser Val Ile Val  
1 5 10 15

&lt;210&gt; 245

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Human cytomegalovirus

&lt;400&gt; 245

Pro Ala Thr Ile Pro Leu Ser Ser Val Ile Val Ala Glu Asn Ser  
1 5 10 15

<210> 246  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 246

Pro Leu Ser Ser Val Ile Val Ala Glu Asn Ser Asp Gln Glu Glu  
1 5 10 15

<210> 247  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 247

Val Ile Val Ala Glu Asn Ser Asp Gln Glu Glu Ser Glu Gln Ser  
1 5 10 15

<210> 248  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 248

Glu Asn Ser Asp Gln Glu Glu Ser Glu Gln Ser Asp Glu Glu Glu  
1 5 10 15

<210> 249  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 249

Gln Glu Glu Ser Glu Gln Ser Asp Glu Glu Glu Glu Gly Ala  
1 5 10 15

<210> 250  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 250

Glu Gln Ser Asp Glu Glu Glu Glu Gly Ala Gln Glu Glu Arg

1 5 10 15

<210> 251  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 251

Glu Glu Glu Glu Glu Gly Ala Gln Glu Glu Arg Glu Asp Thr Val  
1 5 10 15

<210> 252  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 252

Glu Gly Ala Gln Glu Glu Arg Glu Asp Thr Val Ser Val Lys Ser  
1 5 10 15

<210> 253  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 253

Glu Glu Arg Glu Asp Thr Val Ser Val Lys Ser Glu Pro Val Ser  
1 5 10 15

<210> 254  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 254

Asp Thr Val Ser Val Lys Ser Glu Pro Val Ser Glu Ile Glu Glu  
1 5 10 15

<210> 255  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 255

Val Lys Ser Glu Pro Val Ser Glu Ile Glu Glu Val Ala Pro Glu  
1 5 10 15

<210> 256  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 256

Pro Val Ser Glu Ile Glu Glu Val Ala Pro Glu Glu Glu Glu Asp  
1 5 10 15

<210> 257  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 257

Ile Glu Glu Val Ala Pro Glu Glu Glu Glu Asp Gly Ala Glu Glu  
1 5 10 15

<210> 258  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 258

Ala Pro Glu Glu Glu Glu Asp Gly Ala Glu Glu Pro Thr Ala Ser  
1 5 10 15

<210> 259  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 259

Glu Glu Asp Gly Ala Glu Glu Pro Thr Ala Ser Gly Gly Lys Ser  
1 5 10 15

<210> 260  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 260

Ala Glu Glu Pro Thr Ala Ser Gly Gly Lys Ser Thr His Pro Met  
1 5 10 15

<210> 261

<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 261

Thr Ala Ser Gly Gly Lys Ser Thr His Pro Met Val Thr Arg Ser  
1 5 10 15

<210> 262  
<211> 15  
<212> PRT  
<213> Human cytomegalovirus

<400> 262

Gly Lys Ser Thr His Pro Met Val Thr Arg Ser Lys Ala Asp Gln  
1 5 10 15

**THIS PAGE BLANK (USPTO)**